

Colchicine-like β -acetamidoketones as inhibitors of microtubule polymerization: Design, synthesis and biological evaluation of *in vitro* anticancer activity

Ehsan Karimikia^{1,2}, Javad Behravan¹, Afshin Zarghi³, Morteza Ghandadi², Sina Omid Malayeri², Razieh Ghodsi^{1,2*}

¹ Biotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran

² Department of Medicinal Chemistry, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

³ Department of Pharmaceutical Chemistry, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article type:
Original article

Article history:
Received: Sep 9, 2018
Accepted: Apr 7, 2019

Keywords:
Anticancer activity
Chalcone
Colchicine
Colchicine-like
Molecular docking
Tubulin polymerization
 β -acetamidoketones

ABSTRACT

Objective(s): In this study a series of novel colchicine-like β -acetamidoketones was designed and synthesized as potential tubulin inhibitors

Materials and Methods: The cytotoxicity of the novel synthesized β -acetamidoketones was assessed against two cancerous cell lines including MCF-7 (human breast cancer cells) and A549 (adenocarcinomic human alveolar basal epithelial cells) employing the MTT test. Tubulin polymerization test was done by using a commercial kit (tubulin polymerization assay kit).

Results: In general, the cytotoxicity activities were highly dependent on the aromatic substitution pattern of phenyl ring at β position of β -acetamidoketones. Based upon, compound **4f** possessing the same structural elements of colchicine and chalcone 1, revealed the most cytotoxicity more than the other β -acetamidoketone against the cancerous cell lines and showed moderate antitubulin effect. The tubulin inhibitory effect of **4f**, colchicine and chalcone 1 were consistent with their antiproliferative activities. Molecular docking studies of **4f**, into the colchicine-binding site of tubulin exhibited possible mode of interaction between this compound and tubulin.

Conclusion: The structure activity relationship (SAR) data attained showed that the presence of trimethoxy phenyl attached to carbonyl group of β -acetamidoketones and a methoxy group at *para* position of the other ring are essential for cytotoxic activity. In general, the cytotoxicity activities were highly dependent on the aromatic substitution pattern of phenyl ring at β position of β -acetamidoketones.

► Please cite this article as:

Karimikia E, Behravan J, Zarghi A, Ghandadi M, Omid Malayeri S, Ghodsi R. Colchicine-like β -acetamidoketones as inhibitors of microtubule polymerization; Design, synthesis and biological evaluation of *in vitro* anticancer activity. Iran J Basic Med Sci 2019; 22:1138-1146. doi: 10.22038/ijbms.2019.34760.8242

Introduction

Cancer is the reason of one-quarter of all expiries in industrialized countries. Nowadays cancer is the second reason of death in the United States, and is predicted to exceed heart illnesses as the leading cause of death in the next years (1). Therefore; the discovery of anticancer, non-toxic and drug-like active compounds is still an urgent demand.

Microtubules play key role in mitosis and are imperative target for the design of novel anticancer agents. Colchicine (Figure 1) a compound isolated from *Colchicum autumnale* and *Gloriosa superba* was the first drug identified to bind tubulin, and it binds at a specific site called the colchicine domain (2). There are various compounds discovered, which bind tubulins at the colchicine binding site, and hence by inhibiting microtubule polymerization cause cell cycle arrest and, lead to cell death. (3). Combretastatins, separated from the South African tree *Combretum caffrum*, are a series of antimetabolic agents. Combretastatin A-4 (CA-4, Figure 1) is a natural anti-tubulin compound which binds to colchicine's binding site of tubulin and exerts

its antimetabolic effect. Many structural modifications of the combretastatin A-4 molecule, such as variation of substitutions on the A- and B-rings, have been reported (4-6). Numerous Chalcones are also reported as tubulin inhibitors (7, 8). Chalcone 1, a combretastatin-like derivative was known to possess potent anticancer activity and tubulin inhibition effect (7, 9).

Colchicine is identified to have significant anti-mitotic, anti-inflammatory and anti-fibrotic effects. Although colchicine possesses notable *in vitro* antitumor effects, its therapeutic uses have been restricted because of the low bioavailability and high toxicity (10). Though it's unfavorable pharmacological profiles, colchicine is still a lead compound for the discovering of possible antimetabolic drugs. Thus, various analogs of colchicine (11-14) have been synthesized with the goal of discovering novel, valuable drugs with more bioavailability and favorable pharmacological effects. Structure-activity studies (15, 16) suggested that the trimethoxy phenyl ring (A) and the methoxy tropone ring (C) of colchicine include the minimal structural features of the compound needed for its binding to tubulin. On the other hand, a study (17)

*Corresponding author: Razieh Ghodsi. Biotechnology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran. Department of Medicinal Chemistry, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. Tel: +98-51-38823255; Fax: +98-51-38823251; Email: ghodsir@mums.ac.ir




Figure 1. Chemical structures of known tubulin inhibitors (lead compounds) and designed compound

revealed that the B-ring of colchicine plays a key role in the stability of tubulin binding while the A and the C-rings have not significant effect on it and colchicine binds at the α/β interface of tubulin. The B-ring binds on the α -subunit and the A and the C-rings bind on the β -subunit.

In the present study we report the design and synthesis of novel β -acetamidoketones possessing pharmacophoric requirements of anti-tubulins with central β -acetamido ketone bridge (Figure 1). They designed to bear some similarity to colchicine and chalcones. We designed the hybrid compounds, combining ring A and a part of ring B of colchicine and a fragment of chalcone 1 with the same linker length. (Figure 1). They were evaluated for their antiproliferative properties against two human cancer cell lines including MCF-7 and A549. The tubulin inhibitory effect of **4f** (the most cytotoxic compound) was also evaluated. In order to get the better structure activity relationship (SAR) data, we also evaluated the cytotoxicity and tubulin inhibitory properties of chalcone 1.

Experimental

All reagents, chemicals and solvents used in the present study were bought from Merck AG and Aldrich Chemical. Melting points were measured using a Thomas-Hoover capillary apparatus. Infrared spectra were attained by a Perkin Elmer Model 1420 spectrometer (Germany). ^1H NMR and ^{13}C NMR spectra were attained by Bruker FT-300 MHz instrument (Bruker Biosciences, USA). Chloroform-D was used as solvent. Coupling constant (J) values were assessed in hertz (Hz) and spin multiples were given as s (singlet), d (double), t (triplet), q (quartet), m (multiplet). The mass spectra were acquired using a 6410 Agilent LCMS triple quadrupole mass spectrometer (LCMS) with an electrospray ionization (ESI) interface (Japan). Elemental analyses were done on a Cos-Tec model EAS

4010 instrument (Cernusco, Italy) and the results were within $\pm 0.4\%$ of the theoretical values.

General procedure for the synthesis of β -acetamido propiophenones

Substituted benzaldehyde (1.0 mmol) and acetyl chloride (0.5 ml) in acetonitrile (3.0 ml), TFA (0.30 mole %) were added to a solution of an appropriate acetophenone (1.0 mmol), and the mixture was stirred for about 30 min at an ice-water bath, and then permitted to warm to room temperature. upon completion the reaction, the mixture was transferred into ice-water (20.0 ml) saturated NaHCO_3 was added to the mixture to adjust the pH to 7, which resulted to precipitation of the target β -acetamidoketone. The precipitate was filtered, washed with hexane and recrystallized in methanol.

β -acetamido- β -(phenyl)-3,4,5-trimethoxy propiophenone (4a)

Yield, 69%; mp= 101-103 °C: IR (KBr) ν (cm^{-1}): 1658 (C=O), 3266 (NH); ^1H NMR (300 MHz) (CDCl_3) δ (ppm): 1.67 (s, 3H, CH_3), 3.35-3.43 (dd, 1H, CH_2 , J=6.6 & 16.2 Hz), 3.78-3.85 (dd, 1H, CH_2 , J=6.6 & 16.2 Hz), 3.88 (s, 6H, OCH_3), 3.90 (s, 3H, OCH_3), 5.53-5.59 (m, 1H, CH), 6.61-6.63 (d, 1H, NH, J=7.5), 7.21 (s, 2H, 3,4,5-trimethoxyphenyl H_2 & H_6), 7.25-7.36 (m, 5H, phenyl); ^{13}C NMR (75 MHz) (CDCl_3) δ (ppm): 23.38, 43.34, 50.48, 56.29, 60.94, 105.69, 126.58, 127.57, 128.71, 131.79, 140.78, 142.85, 153.07, 169.57, 197.14; MS (ESI) m/z: 358.2 [M+1], 380.2 [M+23]. Anal. Calcd for $\text{C}_{20}\text{H}_{23}\text{NO}_5$: C, 67.21; H, 6.49; N, 3.92. Found: C, 67.43; H, 6.31; N, 4.12.

β -acetamido- β -(4-methoxyphenyl)-3,4,5-trimethoxy propiophenone (4b)

Yield, 64%; mp= 118-120°C: IR (KBr) ν (cm^{-1}): 1647 (C=O), 3319 (NH); ^1H NMR (300 MHz) (CDCl_3) δ (ppm): 2.05 (s, 3H, CH_3), 2.33 (s, 3H, CH_3), 3.32-3.34 (dd, 1H, CH_2 , J=6.9 & 16.2 Hz), 3.77-3.84 (dd, 1H, CH_2 , J=6.9

&16.2 Hz), 3.77 (s, 6H, OCH₃), 3.79 (s, 3H, OCH₃), 5.50-5.52 (m, 1H, CH), 6.51-6.53 (d, 1H, NH, J=7.5), 7.14-7.16 (d, 2H, 4-methylphenyl H₃ & H₅, J=7.8), 7.22-7.25 (m, 4H, 3,4,5-trimethoxyphenyl H₂ & H₆ & 4-methylphenyl H₂ & H₆): ¹³CNMR (75 MHz) (CDCl₃) δ (ppm): 21.05, 23.46, 43.37, 50.39, 56.30, 60, 95, 105.72, 126.50, 129.42, 131.81, 137.31, 137.64, 142.83, 153.08, 169.48, 197.26; MS (ESI) m/z: 372.2 [M+1], 394.2 [M+23]. Anal. Calcd for C₂₁H₂₅NO₅: C, 67.91; H, 6.78; N, 3.77. Found: C, 67.63; H, 6.41; N, 3.42.

β -acetamido- β -(4-acetamidophenyl)-3,4,5-trimethoxy propiophenone (4c)

Yield, 69%; mp= 155-157°C: IR (KBr) ν (cm⁻¹): 1654 (C=O), 3061, 3288 (NH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.06 (s, 3H, CH₃), 2.18 (s, 3H, CH₃), 3.32-3.40 (dd, 1H, CH₂, J=6.6 & 16.5 Hz), 3.75-3.82 (dd, 1H, CH₂, J=6.6 & 16.5 Hz), 3.92 (s, 6H, OCH₃), 3.94 (s, 3H, OCH₃), 5.51-5.55 (dd, 1H, CH, J=6.9 & 12.3 Hz), 6.58-6.61 (d, 1H, NH, J=7.8), 7.20 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆), 7.29-7.31 (d, 2H, 4-acetamidophenyl H₃ & H₅, J=8.4), 7.44-7.47 (d, 2H, 4-acetamido phenyl H₂ & H₆, J=8.4); ¹³CNMR (75 MHz) (CDCl₃) δ (ppm): 27.86, 28.93, 49.09, 54.79, 61.03, 65.52, 110.45, 124.64, 132.01, 136.75, 141.25, 143.01, 147.17, 157.75, 173.60, 174.34, 201.13; MS (ESI) m/z: 415.2 [M+1], 437.2 [M+23]. Anal. Calcd for C₂₂H₂₆N₂O₆: C, 67.91; H, 6.78; N, 3.77. Found: C, 67.55; H, 6.91; N, 3.27.

β -acetamido- β -(4-acetoxyphenyl)-3,4,5-trimethoxy propiophenone (4d)

Yield, 55%; mp= 137-140 °C: IR (KBr) ν (cm⁻¹): 1668, 1751 (C=O), 3284 (NH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.00 (s, 3H, CH₃), 2.28 (s, 3H, CH₃), 3.32-3.39 (dd, 1H, CH₂, J=6.6 & 16.5 Hz), 3.70-3.77 (dd, 1H, CH₂, J=6.6 & 16.5 Hz), 3.90 (s, 6H, OCH₃), 3.91 (s, 3H, OCH₃), 5.49-5.55 (m, 1H, CH), 6.75-6.78 (d, 1H, NH, J=7.5), 7.19 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆), 7.01-7.04 (d, 2H, 4-acetoxyphenyl H₃ & H₅, J=8.4), 7.33-7.36 (d, 2H, 4-acetoxyphenyl H₂ & H₆, J=8.4); ¹³CNMR (75 MHz) (CDCl₃) δ (ppm): 21.11, 23.35, 43.20, 49.91, 56.30, 60.95, 105.67, 121.78, 127.82, 131.67, 138.39, 142.92, 149.90, 153.10, 169.49, 169.65, 196.95; MS (ESI) m/z: 416.2 [M+1], 438.2 [M+23]. Anal. Calcd for C₂₂H₂₅NO₇: C, 67.91; H, 6.78; N, 3.77. Found: C, 68.15; H, 6.41; N, 3.67.

β -acetamido- β -(3-acetoxy, 4-methoxyphenyl)-3,4,5-trimethoxy propiophenone (4e)

Yield, 56%; mp= 183-185 °C: IR (KBr) ν (cm⁻¹): 1618, 1664, 1758 (C=O), 3240 (NH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.03 (s, 3H, CH₃), 2.31 (s, 3H, CH₃), 3.33-3.39 (dd, 1H, CH₂, J=6.9 & 16.5 Hz), 3.76-3.81 (m, 4H, CH₂ & OCH₃), 3.82-3.85 (m, 9H, OCH₃), 5.46-5.51 (m, 1H, CH), 6.50-6.52 (d, 1H, NH, J=7.6 Hz), 6.91-6.93 (d, 1H, 3-acetoxy-4-methoxyphenyl H₅, J=8.4 Hz), 7.05-7.06 (d, 1H, 3-acetoxy-4-methoxyphenyl H₂, J=2.4 Hz), 7.18-7.20 (dd, 1H, 3-acetoxy-4-methoxyphenyl H₆, J= 2.4 & 8.4 Hz), 7.22 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆); ¹³CNMR (75 MHz) (CDCl₃) δ (ppm): 20.64, 23.32, 43.21, 49.73, 55.91, 56.29, 60.92, 105.68, 105.80, 112.36, 121.38, 125.21, 131.71, 133.31, 139.70, 142.81, 150.42, 153.07, 168.94, 169.63, 196.95; MS (ESI) m/z: 446.2 [M+1], 468.2 [M+23]. Anal. Calcd for C₂₃H₂₇NO₈: C, 67.91; H,

6.78; N, 3.77. Found: C, 68.25; H, 6.81; N, 3.68

β -acetamido- β -(4-acetoxy, 3-methoxyphenyl)-3,4,5-trimethoxy propiophenone (4g)

Yield, 67%; mp= 135-137 °C: IR (KBr) ν (cm⁻¹): 1617, 1774 (C=O), 3240 (NH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.01 (s, 3H, CH₃), 2.28 (s, 3H, CH₃), 3.31-3.35 (dd, 1H, CH₂, J=6.9 & 16.41 Hz), 3.74-3.79 (dd, 1H, CH₂, J=6.9 & 16.41 Hz), 3.80 (s, 3H, OCH₃), 3.88-4.04 (m, 9H, OCH₃), 5.44-5.48 (m, 1H, CH), 6.57 (s, 1H, NH), 6.88-6.90 (d, 1H, 4-acetoxy-3-methoxyphenyl H₅, J=8.51), 7.02-7.03 (d, 1H, 4-acetoxy-3-methoxyphenyl H₂, J=2.12 Hz), 7.15-7.17 (dd, 1H, 4-acetoxy-3-methoxyphenyl H₆, J= 2.1 & 8.51 Hz), 7.20 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆); ¹³CNMR (75 MHz) (CDCl₃) δ (ppm): 20.66, 23.43, 43.08, 49.78, 55.94, 56.33, 60.95, 105.72, 112.40, 121.40, 125.21, 131.71, 133.24, 139.75, 142.92, 150.48, 153.12, 168.93, 169.50, 197.05; MS (ESI) m/z: 446.2 [M+1], 468.2 [M+23]. Anal. Calcd for C₂₃H₂₇NO₈: C, 67.91; H, 6.78; N, 3.77. Found: C, 68.29; H, 6.86; N, 3.88.

β -acetamido- β -(3,4-dimethoxyphenyl)-3,4,5-trimethoxy propiophenone (4h)

Yield, 81%; mp= 125-127 °C: IR (KBr) ν (cm⁻¹): 1653 (C=O), 3316 (NH); ¹H NMR (400 MHz) (CDCl₃) δ (ppm): 2.06 (s, 3H, CH₃), 3.31-3.38 (dd, 1H, CH₂, J=7.2 & 16.2 Hz), 3.77-3.84 (dd, 1H, CH₂, J=7.2 & 16.2 Hz), 3.86 (s, 3H, OCH₃), 3.88 (s, 3H, OCH₃), 3.93 (s, 6H, OCH₃), 3.94 (s, 3H, OCH₃), 5.47-5.50 (m, 1H, CH), 6.48-6.50 (d, 1H, NH, J=7.8), 6.81-6.90 (m, 3H, 4-dimethoxyphenyl H₂, H₅ & H₆), 7.23 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆); ¹³C NMR (75 MHz) (CDCl₃) δ (ppm): 23.47, 43.36, 50.59, 55.88, 55.92, 56.31, 60.95, 105.74, 110.44, 111.20, 118.58, 131.81, 133.17, 142.88, 148.49, 149.05, 153.10, 169.46, 197.33; MS (ESI) m/z: 418.2 [M+1], 440.2 [M+23]. Anal. Calcd for C₂₂H₂₇NO₇: C, 67.91; H, 6.78; N, 3.77. Found: C, 67.59; H, 6.86; N, 3.98.

β -acetamido- β -(3,4,5-trimethoxyphenyl)-3,4,5-trimethoxy propiophenone (4i)

Yield, 71%; mp= 133-135 °C: IR (KBr) ν (cm⁻¹): 1659 (C=O), 3246 (NH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 1.74 (s, 3H, CH₃), 3.28-3.35 (dd, 1H, CH₂, J=6.9 & 15.9 Hz), 3.72-3.79 (dd, 1H, CH₂, J=6.9 & 15.9 Hz), 3.81 (s, 3H, OCH₃), 3.83 (s, 6H, OCH₃), 3.92 (s, 6H, OCH₃), 3.95 (s, 3H, OCH₃), 5.44-5.46 (m, 1H, CH), 6.55 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆), 6.60-6.62 (d, 1H, NH, J=7.5), 7.20 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆); ¹³C NMR (75 MHz) (CDCl₃) δ (ppm): 23.36, 43.40, 51.01, 56.09, 56.29, 60.73, 60.93, 103.88, 105.71, 131.82, 136.53, 137.29, 142.86, 153.08, 153.29, 169.54, 197.22; MS (ESI) m/z: 448.2 [M+1], 470.2 [M+23]. Anal. Calcd for C₂₃H₂₉NO₈: C, 61.73; H, 6.53; N, 3.13. Found: C, 62.19; H, 6.76; N, 3.26.

β -acetamido- β -(3-acetoxy, 4-methoxyphenyl)-4-methoxy propiophenone (4j)

Yield, 65%; mp= 156-157°C: IR (KBr) ν (cm⁻¹): 1648, 1691, 1765 (C=O), 3281 (NH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.01 (s, 3H, CH₃), 2.30 (s, 3H, CH₃), 3.37-3.43 (dd, 1H, CH₂, J=6.4 & 16.4 Hz), 3.66-3.72 (dd, 1H, CH₂, J=6.4 & 16.4 Hz), 3.80 (s, 3H, OCH₃), 5.50-5.55 (m, 1H, CH), 6.76-6.77 (d, 1H, NH, J=7.5 Hz), 6.88-6.90 (d, 1H, 3-acetoxy-4-methoxyphenyl H₅, J= 8.4 Hz), 6.92-6.94

(d, 2H, 4-methoxyphenyl H₃ & H₅, J= 8.8 Hz), 7.05(d, 1H, 3-acetoxy-4-methoxyphenyl H₂, J= 2 Hz), 7.16-7.18 (dd, 1H, 3-acetoxy-4-methoxyphenyl H₆, J= 2 & 8.4 Hz) 7.90-7.92 (m, 2H, 2H, 4-methoxyphenyl H₂&H₆, J= 8.8 Hz); ¹³CNMR (75 MHz) (CDCl₃) δ (ppm): 20.65, 23.40, 42.73, 49.33, 55.51, 55.93, 111.82, 112.33, 113.87, 121.29, 125.04, 129.70, 130.52, 133.88, 139.70, 150.29, 163.83, 169.45, 196.93; MS (ESI) m/z:408.2 [M+23]. Anal. Calcd for C₂₁H₂₃NO₆: C, 65.44; H, 6.02; N, 3.63. Found: C, 65.56; H, 6.36; N, 3.39.

β -acetamido- β -(3-acetoxy, 4-methoxyphenyl)-4-fluoro propiophenone (4l)

Yield, 89%; mp= 155-157°C: IR (KBr) ν (cm⁻¹): 1648, 1691, 1765 (C=O), 3281(NH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.01 (s, 3H, CH₃), 2.30 (s, 3H, CH₃), 3.33-3.39 (dd, 1H, CH₂, J=6.0 & 16.4 Hz), 3.64-3.69 (dd, 1H, CH₂, J=6.0 & 16.4 Hz), 3.80 (s, 3H, OCH₃), 5.48-5.52 (m, 1H, CH), 6.75-6.77 (d, 1H, NH, J=7.5Hz), 6.88-6.90 (dd, 1H, 3-acetoxy-4-methoxyphenyl, H₆, J= 2 & 8 Hz), 6.95-6.98 (m, 2H, 3-acetoxy-4-methoxyphenyl H₂&H₅), 7.10-7.17 (m, 2H, 4-fluorophenyl, H₃ & H₅); MS (ESI) m/z: 374.1 [M+1], 396.1 [M+23]. Anal. Calcd for C₂₀H₂₀FNO₅: C, 64.34; H, 5.40; N, 3.75. Found: C, 64.76; H, 6.27; N, 3.84.

β -acetamido- β -(3,4,5-trimethoxyphenyl)- 4-fluoro propiophenone (4m)

Yield, 85%; mp= 186-188 °C: IR (KBr) ν (cm⁻¹): 1618, 1690 (C=O), 3242 (NH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.08 (s, 3H, CH₃), 3.35-3.41 (dd, 1H, CH₂, J=6.4 & 16.4 Hz), 3.69-3.74 (dd, 1H, CH₂, J=6.4 & 16.4 Hz), 3.81 (s, 3H, OCH₃), 3.83 (s, 6H, OCH₃), 5.44-5.49 (m, 1H, CH) 6.55 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆), 6.67-6.69(d, 1H, NH, J=8Hz), 7.12-7.17(t, 2H, 4-fluorophenyl H₂ & H₆, J=8.8Hz), 7.95-7.98 (m, 2H, 4-fluorophenyl H₃ & H₅); ¹³CNMR (75 MHz) (CDCl₃) δ (ppm): 23.40, 43.33, 50.63, 56.14, 60.77, 103.88, 115.74, 116.03, 130.80, 130.93, 133.11, 136.57, 153.34, 169.62, 197.04; MS (ESI) m/z: 376.2[M+1], 398.1 [M+23].

General procedure for deacetylation of acetoxy β -acetamido ketones

A mixture of acetoxy β -acetamido ketones (2.5 mmol) and K₂CO₃ (4 mmol) in NMP (2.5 ml) were heated at 100°C. After completion of the reaction (monitored by TLC), permitted to warm to room temperature and diluted with 2% aqueous NaOH (10 ml) and extracted with diethyl ether. After acidifying the aqueous layer with HCl (6 M) and extracting with diethyl ether, the ethereal phase was dehydrated using Na₂SO₄ and vaporized under vacuum to obtain the target hydroxylated β -acetamido ketones.

β -acetamido- β -(3-hydroxy, 4-methoxyphenyl)-3,4,5-trimethoxy propiophenone (4f)

Yield, 82%; mp= 123-125 °C: IR (KBr) ν (cm⁻¹): 1659 (C=O), 3303 (NH, OH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.05 (s, 3H, CH₃), 3.28-3.36 (dd, 1H, CH₂, J=6.9 & 15.9 Hz), 3.74-3.81(dd, 1H, CH₂, J=6.9 & 15.9 Hz), 3.87 (s, 3H, OCH₃), 3.88-4.01 (m, 9H, OCH₃), 5.45-5.46 (m, 1H, CH), 5.73 (s, 1H, OH), 6.51-6.53 (d, 1H, NH, J=6.9), 6.80-6.92(m, 3H,3-hydroxy-4-methoxyphenyl H₂,

H₃&H₆) 7.22 (s, 2H, 3,4,5-trimethoxyphenyl H₂ & H₆); MS (ESI) m/z:404.2 [M+1], 426.2 [M+23]. Anal. Calcd for C₂₁H₂₅NO₇: C, 62.52; H, 6.25; N, 3.47. Found: C, 62.87; H, 6.44; N, 3.73.

β -acetamido- β -(3-hydroxy, 4-methoxyphenyl)-4-methoxy propiophenone (4k)

Yield, 84%; mp= 120-122 °C: IR (KBr) ν (cm⁻¹): 1664 (C=O), 3357(NH, OH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 2.03 (s, 3H, CH₃), 3.30-3.38 (dd, 1H, CH₂, J=6.0&16.5 Hz), 3.63-3.70 (dd, 1H, CH₂, J=6.0&16.5 Hz), 3.85 (s, 3H, OCH₃), 3.88 (s, 3H, OCH₃), 5.45-5.48 (m, 1H, CH), 5.71 (s, 1H, NH), 6.72 (s, 1H, OH), 6.75-6.95(m, 5H,3-hydroxy-4-methoxyphenyl H₂, H₅ & H₆ & 4-methoxyphenyl H₃ & H₅), 7.91-7.94 (d, 2H, 4-methoxyphenyl H₂ & H₆, J=9 Hz); MS (ESI) m/z:709.3 [2M+23]. Anal. Calcd for C₁₉H₂₁NO₅: C, 66.46; H, 6.16; N, 4.08. Found: C, 66.88; H, 6.73; N, 3.97.

3-(3-hydroxy-4-methoxyphenyl)-1-(3,4,5-trimethoxyphenyl)prop-2-en-1-one(Chalcone1)

The same as reference (9): mp= 144-146 °C; IR (KBr) ν (cm⁻¹): 1675 (C=O), 3416(OH); ¹H NMR (300 MHz) (CDCl₃) δ (ppm): 3.93 (s, 3H, OCH₃), 3.94 (s, 9H, OCH₃), 5.72 (s, 1H, OH), 6.87 (d, 1H, 3-hydroxy-4-methoxyphenyl H₅, J= 8.0 Hz), 7.13 (dd, 1H, 3-hydroxy-4-methoxyphenyl H₆, J= 2.0, 8.0 Hz), 7.25-7.30 (m, 3H, 3-hydroxy-4-methoxyphenyl H₂ & 3,4,5-trimethoxyphenyl, H₂&H₆), 7.33-7.36 (d, 1H, CH, J= 15.5 Hz), 7.72-7.7536 (d, 1H, CH, J= 15.5 Hz); MS (ESI) m/z: 345.1 [M+1], 367.1 [M+23]. Anal. Calcd for C₁₉H₂₀O₆: C, 66.27; H, 5.85. Found: C, 66.58; H, 5.94.


Cytotoxicity assay

General procedure

The MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide) based assay was performed by seeding 5000 cancerous cells per 180 μ l RPMI complete culture medium in each well of 96-well culture plates (18-22). After 24 hr, culture medium was substituted with medium having positive control colchicine and chalcone 1 as well as diverse concentrations of new synthesized compounds and RPMI as negative control. Then cells were incubated at 37 °C in 5% CO₂ incubator for 48 hr. Then 25 μ l of MTT solution (4 mg ml⁻¹) were added to each well and further incubated at 37 °C for 3 hr, then, formazan crystals were dissolved in DMSO (100 μ l) and plates were read using a plate reader (Synergy H4, USA) at 540 nm. This test was done in triplicate determination each time.

Tubulin polymerization assay

Tubulin polymerization test was done by using a commercial kit (Tubulin Polymerization Assay Kit (Porcine tubulin and Fluorescence based Kit, Cat No BK011P, Cytoskeleton, USA)), based on the manufacturer's procedure (23-25). Tubulin protein was added to tubulin buffer (80 mM PIPES, 2 mM MgCl₂, 0.5 mM EGTA, 1 mM GTP, 60% (v/v) glycerol, pH 6.9) then poured to wells of a 96-well plate possessing the cytotoxic chemicals or vehicle and then mixed well. The effects of compound 4f and Chalcone 1 at 50 and 100 μ M concentrations on tubulin polymerization were investigated.



Scheme 1. Reagents and conditions: (a) Cerium (IV) sulfate, CH_3CN (b) H_3BO_3 , CH_3CN , CH_3COCl (c) TFA, CH_3CN , CH_3COCl (d) K_2CO_3 , NMP

Tubulin polymerization was monitored by detecting the enhancement of the fluorescence because of the addition of a fluorescence reporter into microtubules as polymerization occurs. Polymerization was measured by excitation at 360 nm and emission at 420 nm for 1 hr at 1 min breaks by a plate reader (Synergy H4, USA). Colchicine at 50 and 100 μM concentrations and paclitaxel at 3 μM concentration were used as positive destabilizing and stabilizing controls, respectively.

Molecular modeling

Mode of interaction between **4f** and tubulin was evaluated by docking. 2D structure of the compound was organized in Chem Draw Ultra 8.0 software and 3D structure was prepared by Hyperchem 7 software via molecular mechanic force field pre-optimization followed by AM1 semiempirical calculation. The X-ray crystal structure of tubulin (PDB ID: 1SA0) was copied from the Protein Data Bank (www.rcsb.org). Further changes such as polar hydrogen adding and water molecules deletion was done by MOE software. Compound **4f** was docked into the binding site of tubulin using MOE software. Every atom in a 5 Å about the co-crystallized ligand in crystal coordinates of tubulin was selected as binding site. The docking simulations were performed using triangle matcher placement algorithm in combination with London dG scoring function and force field as refinement process. The top-score docking poses were selected for final ligand–target interaction analysis using LigX module in MOE Software. Validation of docking method was first assessed by docking of co-crystallized ligand into the tubulin binding site (26).

Results

Synthesis


As depicted in scheme 1, In order to synthesize **4f**, at first we let 3,4,5-trimethoxy acetophenone **2a** and 3-hydroxy-4-methoxy benzaldehyde **3e** condense in

acetonitrile in the presence of catalytic amount of Cerium (IV) sulfate according to the reported procedure (27) but no reaction was happened. To get the target **4f**, we examined another reported method (28) employing boric acid as a catalyst and acetyl chloride in acetonitrile. Interestingly, the IR and $^1\text{H-NMR}$ spectra revealed that the chalcone **1** was formed instead. Finally, we tried to do the reaction using trifluoroacetic acid as a catalyst in the presence of acetyl chloride in acetonitrile (29). The presence of a peak in 1758 cm^{-1} in the IR spectrum and also a singlet peak in 2.31 ppm in $^1\text{HNMR}$ spectrum proved the formation of **4e** which possesses the acetoxy group instead of hydroxyl group. Hydrolysis of acetoxy group of **4e** in the presence of potassium carbonate in NMP (N-methyl-2-pyrrolidone) at $100\text{ }^\circ\text{C}$ (28) led to the formation of **4f**. We found that TFA is the best catalyst for synthesis of our β -acetamidoketones (Scheme 2), and acetylation of hydroxyl group was observed while using substituted benzaldehyde bearing hydroxyl group (**3d**, **3e** and **3f**). The compounds were characterized by nuclear magnetic resonance, infrared, mass spectrometry and elementary analysis.

Biological evaluation

In vitro anticancer activity

The cytotoxicity of the synthesized compounds was assessed against two cancerous cell lines including MCF-7 (human breast cancer cells) and A549 (adenocarcinomic human alveolar basal epithelial cells) using the MTT test. As shown in Table 1, by comparing the cytotoxicity of **4j** and **4k** (which displayed no activity at the concentrations below 100 μM), with those of **4e** and **4f**, we can conclude that trimethoxy phenyl moiety attached to carbonyl group is essential for cytotoxic activity of these compounds. The cytotoxicity activity was highly dependent on the aromatic substitution pattern of phenyl ring at β position of β -acetamidoketones. SAR data showed that, introducing an electron donating



Scheme 2. Reagents and conditions: (a) TFA, CH₃CN, CH₃COCl (b) K₂CO₃, NMP, 100 °C

group in the *para* position of **4a** increased its cytotoxic activity against MCF-7 cells (see the anti-proliferative activities of **4a-4d** in Table 1). According to our results, compound **4f** was at least twice more cytotoxic than acetoxy analogue **4e**. Comparing the cytotoxic activity of **4f** with its parent compound **4e** shows that the hydroxyl group at *meta* position of phenyl ring has a crucial role for its activity which may explain the ability of hydroxyl group to form hydrogen bond within the active site of tubulin as a hydrogen binding donor. Replacement of the acetamido group at *para* position of phenyl ring (**4c**) by acetoxy substituent (**4d**) increased cytotoxic activity considerably. Our results indicated that compound **4i** possessing trimethoxy phenyl ring at β position showed more cytotoxicity activities than **4h** having dimethoxy phenyl ring in MCF-7 cells. According

to our results, compound **4f** possessing the colchicine and chalcone 1 aromatic substitution pattern showed the highest cytotoxicity among the β -acetamidoketones series against the cancer cell lines. As chalcone 1 showed higher anti-proliferative activity compared to colchicine, it can be concluded that α , β -unsaturated bridge of chalcone 1 might act as the Michael acceptor in tubulin binding site or other targets in addition to rigid linkage. In general, our compounds exhibited more cytotoxicity effects in MCF-7 cancer cells compared to A549 cancer cells, demonstrating that the compounds may exert their cytotoxicity with different mechanisms in different cancer cells.

Tubulin polymerization assay

In order to clarify whether the cytotoxic activity of **4f**

Table 1. The *in vitro* anti-proliferative activities of colchicine-like β -acetamidoketones, colchicine, and chalcone 1 against human cancer cell lines

Compound	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	MCF-7 IC ₅₀ (μ M)	A549 IC ₅₀ (μ M)
4a	OCH ₃	OCH ₃	OCH ₃	H	H	H	>100	>100
4b	OCH ₃	OCH ₃	OCH ₃	H	CH ₃	H	41.9 \pm 4.72	>100
4c	OCH ₃	OCH ₃	OCH ₃	H	NHCOCH ₃	H	69.7 \pm 9.1	>100
4d	OCH ₃	OCH ₃	OCH ₃	H	OCOCH ₃	H	30.4 \pm 2.78	48.9 \pm 3.98
4e	OCH ₃	OCH ₃	OCH ₃	OCOCH ₃	OCH ₃	H	34.1 \pm 3.5	48.8 \pm 4.2
4f	OCH ₃	OCH ₃	OCH ₃	OH	OCH ₃	H	14.6 \pm 2.1	23.2 \pm 1.95
4g	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCOCH ₃	H	36.2 \pm 3.5	49.3 \pm 3.98
4h	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃	H	37.2 \pm 4.12	37.7 \pm 3.05
4i	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃	18.8 \pm 1.1	40.8 \pm 2.7
4j	H	OCH ₃	H	OCOCH ₃	OCH ₃	H	>100	>100
4k	H	OCH ₃	H	OH	OCH ₃	H	>100	87 \pm 7.5
4l	H	F	H	OCOCH ₃	OCH ₃	H	>100	>100
4m	H	F	H	OCH ₃	OCH ₃	OCH ₃	>100	>100
chalcone 1	-	-	-	-	-	-	0.9 \pm 0.27	2.9 \pm 0.43
colchicine	-	-	-	-	-	-	1.1 \pm 0.37	ND




Figure 2. Effect of compounds (**4f**, Colchicine, Chalcone 1 and Paclitaxel) on *in vitro* tubulin polymerization

was related to the tubulin binding ability, compounds **4f** (the most cytotoxic compound), chalcone 1 and reference compound colchicine (polymerization suppressor) and a polymerization promoter (paclitaxel) were assessed for its effect on tubulin polymerization. As illustrated in Figure 2, **4f** proved to be inhibitor of tubulin polymerization in a manner similar to that of chalcone 1 and colchicine. However, **4f** was not as potent as these reference tubulin inhibitors but its tubulin inhibition effect was dose dependent. SAR acquired data indicated that flexible bridge in **4f**, could lead to conformational flexibility that is unfavorable for tubulin inhibition. Because of higher tubulin inhibitory effect of chalcone 1 compared with colchicine it can be concluded that α , β -unsaturated bridge of chalcone might act as the Michael acceptor in tubulin binding site in addition to the rigid linkage. The tubulin inhibitory effect of **4f**, colchicine and chalcone 1 were consistent with their antiproliferative activities. On the other hand, the order of tubulin inhibitory effect of these three compounds is the same as their antiproliferative activities.

Molecular modeling (docking) studies

In an attempt to elucidate the possible binding mode of this novel series of tubulin inhibitors, docking of the most potent compound **4f** was performed at the active site of the tubulin dimer. The quality and validity of docking procedure were investigated by docking of co-crystallized ligand within the binding site of tubulin. From docking studies, the top binding position exhibited a similar alignment in the binding pocket to the co-crystallized ligand found in crystal assembly (PDB ID of 1SA0). The root mean square deviation (RMSD) between co-crystallized ligand within the binding site and ligand docked in the crystal structure of tubulin was 1.4 Å, indicative of a proper ability to mimic the ligand binding mode recognized in the experimental data (30, 31). As mentioned above, compound **4f** possessing structural elements of colchicine and chalcone 1, demonstrated the most anti proliferative activity compared to the other synthesized compounds which inhibited the polymerization of tubulin. According to ligand interaction mode of **4f** by LigX module of MOE software, hydroxyl group of **4f**, could form hydrogen bonds with LysB352 or AlaB317 (Figure 3). The methoxy groups of




Figure 3. The interaction of compound **4f** in the crystal assembly of tubulin

the first phenyl ring made contacts with the backbone of several amino acid residues such as TyrA224, GlyA11, GlnB247 and LysB254. This phenyl ring of **4f** provided additional interaction with LeuB 248. The other ring of **4f** made contacts with the aliphatic chain of residues CysB241, ValB318, AlaB316, AlaB317 and LysB352. The β -acetamido side chain of **4f** was surrounded by residues AlaB354, AlaA180, SerA178, ThrB353, ThrA179 and AsnB249 and could make contacts with them. These hydrophobic interactions and hydrogen bonds formation of **4f** with tubulin binding site can explain inhibitory effect of this compound.

Discussion

A new series of colchicine-like β -acetamidoketone analogues was synthesized and evaluated for their cytotoxic activity against MCF-7 and A549 cancer cells. The structure activity data acquired indicate that the presence of trimethoxy phenyl attached to carbonyl group of and a methoxy group at *para* position of the other ring in the synthesized compounds are essential for cytotoxic activity. As chalcone 1 showed higher antiproliferative activity compared to colchicine, it can be concluded that α , β -unsaturated bridge of chalcone 1 might act as the Michael acceptor in tubulin binding site or other targets in addition to rigid linkage. In general, the cytotoxic activities were highly dependent on the aromatic substitution pattern of the phenyl ring at position β of β -acetamidoketones. Based upon, compound **4f** possessing the trimethoxyphenyl and 3-hydroxy-4-methoxy groups, demonstrated the best cytotoxicity among the other β -acetamidoketones against the cancerous cell lines and proved to be a tubulin inhibitor. Finally, molecular docking studies of **4f** into the colchicine-binding site of tubulin demonstrated the possible interactions of this compound at the active site of tubulin. The hydrophobic interactions and hydrogen bonds formation of **4f** with tubulin binding site can explain inhibitory effect of this compound.

Conclusion

The cytotoxicity activities were highly dependent

on the aromatic substitution pattern of phenyl ring at position β of β -acetamidoketones. Based upon, compound **4f** possessing the same structural elements of colchicine and chalcone **1**, revealed the most cytotoxicity more than the other β -acetamidoketone against the cancerous cell lines and showed moderate antitubulin effect. The tubulin inhibitory effect of **4f**, colchicine and chalcone **1** were consistent with their antiproliferative activities. Molecular docking studies of **4f**, into the colchicine-binding site of tubulin exhibited possible mode of interaction between this compound and tubulin.

Acknowledgment

We are grateful to Research Deputy of Mashhad University of Medical Sciences, Mashhad, Iran for financial support of this research as part of thesis of Ehsan Karimikia.

Conflicts of Interest

The authors declare no conflict of interest.

References

- Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin* 2017; 67: 7-30.
- Chinigo GM, Paige M, Grindrod S, Hamel E, Dakshanamurthy S, Chruszcz M, Minor W, Brown ML. Asymmetric synthesis of 2,3-dihydro-2-arylquinazolin-4-ones: methodology and application to a potent fluorescent tubulin inhibitor with anticancer activity. *J Med Chem* 2008; 51: 4620-4631.
- Abolhasani H, Zarghi A, Abolhasani A, Hamzeh-Mivehroud M, Bargahi N, Notash B, Mojarrad JS, Dastmalchi S. Design, synthesis and *in vitro* cytotoxicity evaluation of new 3',4'-bis (3,4,5-trisubstituted)-4'H-spiro[indene-2,5'-isoxazol]-1(3H)-one derivatives as Promising anticancer agents. *Lett Drug Des Discov* 2014; 11: 1149-1161.
- Hsieh HP, Liou J.P, Mahindroo N. Pharmaceutical design of antimetabolic agents based on combretastatins. *Curr Pharm Des* 2005; 11: 1655-1677.
- Nam NH. Combretastatin A-4 analogues as antimetabolic anticancer agents. *Curr Med Chem* 2003; 10: 1697-1722.
- Tron GC, Pirali T, Sorba G, Pagliai F, Busacca S, Genazzani AA. Medicinal chemistry of combretastatin A4: present and future directions. *J Med Chem* 2006; 49: 3033-3044.
- Kong Y, Wang K, Edler MC, Hamel E, Mooberry SL, Paige MA, Brown M.L. A boronic acid chalcone analog of combretastatin A-4 as a potent anti-proliferation agent. *Bioorg Med Chem* 2010; 18: 971-977.
- Mahapatra DK, Bharti SK, Asati V. Anti-cancer chalcones: Structural and molecular target perspectives. *Eur J Med Chem* 2015; 98: 69-114.
- Ducki S, Rennisona D, Woo M, Kendall A, Fournier J, Chabert D, Gown ATM, Lawrence NJ. Combretastatin-like chalcones as inhibitors of microtubule polymerization. Part 1: synthesis and biological evaluation of antivascular activity. *Bioorg Med Chem* 2009; 17: 7698-7710.
- Kozaka T, Nakagawa-Goto K, Shi Q, Lai CU, Hamel E, Bastow KF, Brossi A, Lee KH. Antitumor agents 273. Design and synthesis of N-alkyl-thiocolchicinoids as potential antitumor agents. *Bioorg Med Chem Lett* 2010; 20: 4091-4094.
- Cifuentes M, Schilling B, Ravindr R, Winterra J, Janik ME. Synthesis and biological evaluation of B-ring modified colchicine and isocolchicine analogs. *Bioorg Med Chem Lett* 2006; 16: 2761-2764.
- Chabert MFD, Vinader V, Santos AR, Redondo-Horcajo M, Dreneau A, Basak R, Cosentino L, Marston G, Abdel-Rahman H, Loadman PM, Shnyder SD, Díaz JF, Barasoain I, Falconer RA, Pors K. Synthesis and biological evaluation of colchicine C-ring analogues tethered with aliphatic linkers suitable for prodrug derivatization. *Bioorg Med Chem Lett* 2012; 22: 7693.
- Boyer FD, Dubois J, Thoret S, Huu Dau MET, Hanna I. Synthesis and tubulin-binding properties of new allicolchicinoids. *Bioorg Chem* 2010; 38: 149-158.
- Nakagawa-Goto K, Chen CX, Hamel E, Wu CC, Bastow KF, Brossi A, Lee KH. Antitumor agents. Part 236: Synthesis of water-soluble colchicine derivatives. *Bioorg Med Chem Lett* 2005; 15: 235.
- Yang B, Zhu ZC, Goodson HV, Miller MJ. Syntheses and biological evaluation of ring-C modified colchicine analogs. *Bioorg Med Chem Lett* 2010; 20: 3831-3833.
- Bhattacharyya B, Panda D, Gupta, S, Banerjee M. Antimitotic activity of colchicine and the structural basis for its interaction with tubulin. *Med Res Rev* 2008; 28: 155-183.
- Chaudhuril AR, Seetharamalu P, Schwarz1 PM, Hausheer FH, Luduena RF. The interaction of the B-ring of colchicine with alpha-tubulin: a novel footprinting approach. *J Mol Biol* 2000; 303: 679-692.
- Malayeri SO, Abnous K, Arab A, Akaberi M, Mehri S, Zarghi A, Ghodsi, R. Design, synthesis and biological evaluation of 7-(aryl)-2,3-dihydro-[1,4]dioxino[2,3-g]quinoline derivatives as potential Hsp90 inhibitors and anticancer agents. *Bioorg Med Chem* 2017; 25:1294-302.
- Ghodsi R, Azizi E, Ferlin MG, Pezzi V, Zarghi A. Design, synthesis and biological Evaluation of 4-(Imidazolylmethyl)-2-Aryl-Quinoline Derivatives as aromatase Inhibitors and anti-breast cancer agents. *Lett Drug Des Discov* 2016; 13:89-97.
- Ghodsi R, Azizi E, Zarghi A. Design, synthesis and biological evaluation of 4-(Imidazolylmethyl)-2-(4-methylsulfonyl phenyl)-quinoline derivatives as selective COX-2 inhibitors and *in vitro* anti-breast cancer agents. *Iran J Pharm Res* 2016; 15:169-77.
- Jafari F, Baghayi H, Lavaee P, Hadizadeh F, Soltani F, Moallemzadeh H, et al. Design, synthesis and biological evaluation of novel benzo- and tetrahydrobenzo-[h]quinoline derivatives as potential DNA-intercalating antitumor agents. *Eur J Med Chem* 2019; 164:292-303.
- Golmakaniyoon S, Askari VR, Abnous K, Zarghi A, Ghodsi R. Synthesis, characterization and in-vitro evaluation of novel naphthoquinone derivatives and related imines: Identification of new anticancer leads. *Iran J Pharm Res* 2019;18:16-29.
- Shobeiri N, Rashedi M, Mosaffa F; Zarghi A, Ghandadi M, Ghasemi A, Ghodsi R. Synthesis and biological evaluation of quinoline analogues of flavones as potential anticancer agents and tubulin polymerization inhibitors. *Eur J Med Chem* 2016; 114:14-23.
- <http://www.cytoskeleton.com/bk011p>
- Bonne D, Heusele C, Simon C, Pantaloni D. 4', 6-Diamidino-2-phenylindole, a fluorescent probe for tubulin and microtubules. *J Biol Chem* 1985; 260: 2819-2825.
- Omid Malayeri S, Tayarani-Najaran Z, Shaebani Behbahani F, Rashidi R, Delpazir S, Ghodsi R. Synthesis and biological evaluation of benzo[b]furo[3,4-e][1,4]diazepin-1-one derivatives as anti-cancer agents. *Bioorg Chem* 2018; 80:631-638.
- Selvam, N.P, Perumal, P.T. Cerium (IV) sulfate catalyzed simple and convenient synthesis of β -acetamidocarbonyl compounds. *ARKIVOC*, 2009; 10: 265-282.
- Karimi-Jaberi, Z, Mohammadi, K. One-pot synthesis of using boric acid at room temperature. *Sci Word J* 2012, Article ID 925617.
- Zhang XH, Fan L, Liu J, Yang DC. One-pot synthesis of

β -acetamido- β -arylpropiophenone employing trifluoroacetic acid as an efficient catalyst. *Res Chem Inter med* 2011; 37:811-820.

29. Chakraborti A, Sharma L, Sharma U. A mild and chemoselective method for deprotection of aryl acetates and benzoates under non-hydrolytic condition. *Tetrahedron*. 2001; 57:9343-9346.

30. Abdizadeh T, Kalani MR, Abnous K, Tayarani-Najaran Z,

Khashyarmanesh BZ, Abdizadeh R, Ghodsi R, Hadizadeh F. Design, synthesis and biological evaluation of novel coumarin-based benzamides as potent histone deacetylase inhibitors and anticancer agents. *Eur J Med Chem* 2017; 132:42-62.

31. Aboutorabzadeh SM, Mosaffa F, Hadizadeh F, Ghodsi R. Design, synthesis, and biological evaluation of 6-methoxy-2-arylquinolines as potential P-glycoprotein inhibitors. *Iran J Basic Med Sci* 2018;21:9-18.