

## Enzyme inhibitory and radical scavenging effects of some antidiabetic plants of Turkey

Nilüfer Orhan<sup>1</sup>, Sanem Hoşbaş<sup>1</sup>, Didem Deliorman Orhan<sup>1\*</sup>, Mustafa Aslan<sup>1</sup>, Fatma Ergun<sup>1</sup>

<sup>1</sup>Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, 06330 Hipodrom, Ankara, Turkey

### ARTICLE INFO

#### Article type:

Original article

#### Article history:

Received: Jul 6, 2013

Accepted: Dec 21, 2013

#### Keywords:

$\alpha$ -amylase  
 $\alpha$ -glucosidase  
*Gentiana olivieri*  
*Helichrysum*  
*Juniperus*  
*Viscum album*

### ABSTRACT

**Objective(s):** Ethnopharmacological field surveys demonstrated that many plants, such as *Gentiana olivieri*, *Helichrysum graveolens*, *Helichrysum plicatum* ssp. *plicatum*, *Juniperus oxycedrus* ssp. *oxycedrus*, *Juniperus communis* var. *saxatilis*, *Viscum album* (ssp. *album*, ssp. *austriacum*), are used as traditional medicine for diabetes in different regions of Anatolia. The present study was designed to evaluate the *in vitro* antidiabetic effects of some selected plants, tested in animal models recently.

**Materials and Methods:**  $\alpha$ -glucosidase and  $\alpha$ -amylase enzyme inhibitory effects of the plant extracts were investigated and Acarbose was used as a reference drug. Additionally, radical scavenging capacities were determined using 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) ABTS radical cation scavenging assay and total phenolic content of the extracts were evaluated using Folin Ciocalteu method.

**Results:** *H. graveolens* ethanol extract exhibited the highest inhibitory activity (55.7 %  $\pm$  2.2) on  $\alpha$ -amylase enzyme. Additionally, *J. oxycedrus* hydro-alcoholic leaf extract had potent  $\alpha$ -amylase inhibitory effect, while the hydro-alcoholic extract of *J. communis* fruit showed the highest  $\alpha$ -glucosidase inhibitory activity (IC<sub>50</sub>: 4.4  $\mu$ g/ml).

**Conclusion:** Results indicated that, antidiabetic effect of hydro-alcoholic extracts of *H. graveolens capitulums*, *J. communis* fruit and *J. oxycedrus* leaf might arise from inhibition of digestive enzymes.

#### ► Please cite this paper as:

Orhan N, Hoşbaş S, Orhan DD, Aslan M, Ergun F. Enzyme inhibitory and radical scavenging effects of some antidiabetic plants of Turkey. Iran J Basic Med Sci. 2014; 17:426-432.

### Introduction

Diabetes mellitus is a growing health problem worldwide causing severe and costly complications including blindness, cardiac and kidney diseases (1). According to Shaw *et al* (2010), the world prevalence of diabetes among adults will increase to 7.7%, and affect 439 million adults by 2030. Between 2010 and 2030, there will be a 69% increase in number of adults with diabetes in developing countries and a 20% increase in developed countries (2).

Approaches to the control of blood glucose and prevention of hyperglycemia are central to the treatment of diabetes mellitus. Appetite suppressants, inhibitors of digestion, insulin secretagogues, insulin potentiators, insulin mimetics, stimulants of glucose utilization, inhibitors of gluconeogenesis and glucogenolysis are used to balance blood glucose. At present, none of these therapies either alone or in combination can redraw normal blood glucose homeostasis. Additionally many limitations exist in the use of anti-diabetic drugs; medicines available for management of diabetes exert serious side effects such as hepatotoxicity, abdominal pain, flatulence, diarrhea, and hypoglycemia. Also after prolonged treatment,

drug resistance is reported for these medicines (3-6). Therefore, researchers have targeted towards the discovery of drug candidates from potential sources. Traditional medicines play an important role as starting material for drug discovery. For documentation of ethnopharmacological knowledge, many comprehensive field surveys have been conducted all over the world for years and many plants used against diabetes have been recorded (7-10).

Antidiabetic activities of plants used against diabetes in Turkey as folk medicine were studied in detail by our research group. In our research on *in vivo* antidiabetic activity of traditional medicines from 2000, seven plant species including *Gentiana olivieri* Griseb (Gentianaceae), *Helichrysum graveolens* (Bieb.) Sweet (Asteraceae), *H. plicatum* ssp. *plicatum* DC. (Asteraceae), *Juniperus oxycedrus* ssp. *oxycedrus* L. (Cupressaceae), *J. communis* var. *saxatilis* Pall. (Cupressaceae), *Viscum album* L. (ssp. *album* and ssp. *austriacum*) (Loranthaceae) were evaluated for their antidiabetic activity (11-16). Due to their promising antidiabetic effect in *in vivo* studies, they were selected as the subject of this study.

\*Corresponding author: Didem Deliorman Orhan. Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, 06330, Etiler, Ankara, Turkey. Tel: +90-312 2023173; Fax: +90-312-2235018; email: didem@gmail.com

**Table 1.** General information about plants used in the study

Plant	Place of collection	Voucher No.	Part used	Extraction method	Extract	Yield % (w/w)
<i>Gentiana olivieri</i> Griseb.	Oğuzeli, Gaziantep	GUE 2621	Aerial part	Decoction	HA	38.7
<i>Helichrysum graveolens</i> (Bieb.) Sweet	Ilgaz Mt., Kastamonu	GUE 2356	Capitulum	Maceration	HA	11.5
<i>H. plicatum</i> ssp. <i>plicatum</i> DC.	Palandöken Mt., Erzurum	GUE 2355	Capitulum	Maceration	HA	19.3
				Infusion	Aq	17.5
<i>Juniperus communis</i> var. <i>saxatilis</i> Pall.	Akdağmadeni, Yozgat	GUE 2617	Fruit	Maceration	HA	36.0
				Leaf	HA	29.0
<i>J. oxycedrus</i> ssp. <i>oxycedrus</i> L.	Akdağmadeni, Yozgat	GUE 2616	Fruit	Maceration	HA	33.3
				Infusion	Aq	26.0
				Leaf	HA	35.2
<i>Viscum album</i> ssp. <i>album</i> L.	Bağlum, Ankara	AEF 18953	Aerial part	Maceration	HA	43.2
				Infusion	Aq	25.9
<i>V. album</i> ssp. <i>austriacum</i> (Wiesb.)	Kızılcahamam, Ankara	AEF 18939	Aerial part	Maceration	HA	41.2
				Infusion	Aq	27.6

AEF: Herbarium of Faculty of Pharmacy at Ankara University, GUE: Herbarium of Faculty of Pharmacy at Gazi University, HA: Hydro-alcoholic, Aq: Aqueous

The plants used in this study are well known and widely consumed as food and medicine in different regions of Anatolia. Aerial parts of *G. olivieri* are used as bitter tonic, appetizer, antidiabetic, antipyretic, stomachic, and for mental disorders. Gentians are also used in small amounts as food and beverage flavoring, in antismoking products and even as a substitute for hops in beer making. *Helichrysum* species have been used as diuretics, lithagogues, anti-asthmatics, for stomachache, and against kidney stones. The capitulums of *Helichrysum* species are used to decrease blood glucose levels and aerial parts are also marketed as herbal tea in herbal stores. Juniper berries and leaves are used for antidiabetic, diuretic, antiseptic, carminative, stomachic, antirheumatic, antifungal, and disinfectant properties in many folk medicines (13, 14). Also, berries are used as spice in European cuisine to impart a sharp, clear flavor to meat dishes, pork, cabbage, and sauerkraut dishes (18). Twigs and leaves of *V. album* (European mistletoe) are used for many therapeutic applications such as diabetes mellitus, chronic cramps, stroke, stomach problems, heart palpitations, hypertension, and breathing difficulties (15). Additionally leaves of *V. album* are used as tea for bracing and fruits are eaten fresh and pickled in Turkey (19).

The goal of the present study is to determine the inhibitory effects of the selected plants that were found to have *in vivo* antidiabetic activity on carbohydrate digestion enzymes such as  $\alpha$ -amylase and  $\alpha$ -glucosidase. Inhibition of these enzymes, involved in the digestion of carbohydrates, can significantly reduce the post-prandial increase of blood glucose. So, plants with inhibitory effects on these enzymes might be beneficial in diabetic patients. Oxidative stress, is one of the major problems observed during hyperglycemia and it contributes to severe complications in diabetics (20). Plants with both antidiabetic and antioxidant effects could be useful for people suffering from diabetes mellitus. Therefore, ABTS radical scavenging activity

and total phenolic contents of the extracts were also determined.

## Materials and Methods

### Plant materials

Plant materials were collected from different localities of Turkey and identified by researchers. Voucher specimens are preserved in the herbariums of Gazi and Ankara Universities, Faculty of Pharmacy, (Ankara), Turkey. Plant names, parts used, collection sites and herbarium numbers of the plants are given in Table 1.

### Preparation of extracts

Aqueous and hydro-alcoholic extracts of the plants were prepared according to folkloric usage as described in the previous *in vivo* antidiabetic activity studies (11–16). For decoctions, 1 g of air-dried plant material was added to 100 ml of distilled water and boiled on slow heat for 30 min. Infusions were prepared by pouring 100 ml of boiling water onto 1 g of dried plant material. The extraction continued for 30 min while cooling. Hydro-alcoholic extracts were prepared by maceration of 1 g of powdered material with 100 ml of ethanol (80%) at room temperature for 8 hr. Extracts were then filtered through filter paper and condensed by a rotary evaporator. Extraction yields were calculated after freeze-drying till dryness. Extract yields and other information are given in Table 1.

### Assay for $\alpha$ -amylase inhibitory activity

The  $\alpha$ -amylase inhibition method was performed using the chromogenic method of Ali *et al* (21). Porcine pancreatic  $\alpha$ -amylase (EC 3.2.1.1, type VI, Sigma) was dissolved in ice-cold distilled water (4 U/ml). As substrate solution, potato starch (0.5 %, w/v) in 20 mM phosphate buffer (pH 6.9) was used. Experiments were carried out with three replicates.

40  $\mu$ l of plant extract in DMSO, 160  $\mu$ l of distilled water and 400  $\mu$ l of starch were mixed in an Eppendorf tube. The reaction was initiated by the

**Table 2.**  $\alpha$ -Amylase inhibitory activity of plant extracts

Material/Plant	Part used	Extract	Inh. % $\pm$ SD ( $\mu$ g/ml)			
			3000	1000	300	100
Acarbose	-	-	73.7 $\pm$ 0.6	67.2 $\pm$ 0.6	51.8 $\pm$ 2.9	32.6 $\pm$ 0.3
<i>Gentiana olivieri</i>	Aerial part	HA	39.6 $\pm$ 0.4	13.9 $\pm$ 0.6	-	-
<i>Helichrysum graveolens</i>	Capitulum	Aq	3.5 $\pm$ 1.8	-	-	-
		HA	55.7 $\pm$ 2.2	15.7 $\pm$ 1.6	-	-
<i>Helichrysum plicatum</i>	Capitulum	Aq	12.7 $\pm$ 2.8	13.7 $\pm$ 1.5	16.7 $\pm$ 1.4	17.5 $\pm$ 0.8
		HA	5.4 $\pm$ 2.3	-	-	-
<i>Juniperus communis</i> var. <i>saxatilis</i>	Leaf	HA	53.6 $\pm$ 0.8	2.4 $\pm$ 2.4	-	-
	Fruit	HA	29.8 $\pm$ 1.2	22.6 $\pm$ 1.7	-	-
<i>J. oxycedrus</i> ssp. <i>oxycedrus</i>	Fruit	Aq	8.2 $\pm$ 6.5	-	-	-
		HA	52.6 $\pm$ 0.8	39.0 $\pm$ 1.0	-	-
<i>Viscum album</i> ssp. <i>album</i>	Aerial part	Aq	42.1 $\pm$ 2.0	11.3 $\pm$ 4.6	-	-
		HA	51.7 $\pm$ 0.9	25.6 $\pm$ 0.9	25.2 $\pm$ 1.3	25.0 $\pm$ 0.7
<i>V. album</i> ssp. <i>austriacum</i>	Aerial part	Aq	14.0 $\pm$ 4.2	2.2 $\pm$ 1.5	-	-
		HA	8.7 $\pm$ 2.3	2.4 $\pm$ 1.2	2.0 $\pm$ 1.8	1.8 $\pm$ 0.6
<i>V. album</i> ssp. <i>austriacum</i>	Aerial part	Aq	-	-	-	9.0 $\pm$ 3.3
		HA	44.3 $\pm$ 4.1	10.8 $\pm$ 3.0	2.6 $\pm$ 2.1	-

n=3, SD: Standard deviation, -:no activity, HA: Hydro-alcoholic, Aq: Aqueous

addition of 200  $\mu$ l of the enzyme solution. The tubes were incubated at 37°C for 5 min. After that, 200  $\mu$ l of this mixture was added into another tube containing 100  $\mu$ l DNS color reagent solution (96 mM 3, 5-dinitrosalicylic acid, 5.31 M sodium potassium tartrate in 2 M NaOH) and put into a 85°C heater. After 15 min, this mixture was diluted with 900  $\mu$ l distilled water and taken from the heater. Tubes were cooled on ice and the absorbance of the mixture was read at 540 nm. Acarbose was used as the positive control. The absorbance (A) due to maltose generated was calculated according to following formula:

$$A_{540\text{nm}} \text{ control or plant extract} = A_{540\text{nm}} \text{ Test} - A_{540\text{nm}} \text{ Blank}$$

The amount of maltose generated was calculated by using the maltose standard calibration curve (0 – 0.1% w/v) and the obtained net absorbance. Percent of inhibition was calculated as:

$$\% \text{ inhibition} = [1 - (\text{mean maltose in sample} / \text{mean maltose in control})] \times 100$$

#### Assay for $\alpha$ -glucosidase inhibitory activity

$\alpha$ -Glucosidase activity was performed according to the method of Lam *et al* (22).  $\alpha$ -Glucosidase type IV enzyme (Sigma Co., St. Louis, USA) from *B. stearothermophilus* was dissolved in 0.5 M phosphate buffer (pH 6.5) (3 U/ml). The enzyme solution (20  $\mu$ l) and test extracts (10  $\mu$ l) dissolved in MeOH-H<sub>2</sub>O (1:9, v/v) were preincubated in a 96-well microtiter plate for 15 min at 37°C. After that, the substrate solution [10  $\mu$ l, 20 mM *p*-nitrophenyl- $\alpha$ -D-glucopyranoside (NPG), Sigma] in the same buffer was added. The mixture was incubated for 35 min at 37°C. The increase in the absorption at 405 nm due to the hydrolysis of NPG by  $\alpha$ -glucosidase was measured by an ELISA microtiter plate reader. Acarbose (Bayer Group, Turkey) was used as a

positive control. The inhibition percentage (%) was calculated by the equation:

$$\text{Inhibition (\%)} = [1 - (A_{\text{sample}} / A_{\text{control}})] \times 100$$

IC<sub>50</sub> calculations were done by using Sigma Plot 12.0 software. Minimum of eight different concentrations prepared from the stock solutions of extracts were used for calculating the IC<sub>50</sub> value. The logarithmic concentrations (10.000–0.1  $\mu$ g/ml) were chosen.

#### Assay for scavenging activity of ABTS radical cation

ABTS radical cation (ABTS<sup>+</sup>) scavenging assay was achieved by using the spectrophotometric methods of Re *et al* (23) and Meot-Duros *et al* (24) with slight modifications. ABTS (7 mM) was dissolved in distilled water and the ABTS radical cation was generated by adding 2.45 mM potassium per-sulfate. The radical production was completed after incubation for 16 hr in the dark at 20°C. Absorbance of ABTS solution was adjusted to 0.7  $\pm$  0.02 at 734 nm by the addition of phosphate buffer solution (PBS) at pH 7.4. 1 ml diluted ABTS solution was added to 10  $\mu$ l of extract (PBS or Trolox). Samples were vortexed and their absorbances were read versus PBS blank at 734 nm. Trolox was used as the positive control. The inhibition percentage was calculated according to the following formula:

$$\text{Inhibition percentage} = [1 - (A_{\text{extract}} / A_{\text{control}})] \times 100$$

#### Determination of total phenol content

The extracts (100  $\mu$ l) were mixed with 0.2 ml Folin-Ciocalteu reagent, 2 ml of H<sub>2</sub>O, and 1 ml of 15 % Na<sub>2</sub>CO<sub>3</sub>, respectively. The absorbance of mixture was measured at 765 nm after 2 hr at room temperature. The mean of three readings was used and the total phenol content was expressed in mg of gallic acid equivalents (GAE)/g extracts (25). The coefficient of determination was r<sup>2</sup> = 0.9957.

**Table 3.**  $\alpha$ -Glucosidase inhibitory activity of plant extracts and total phenol content (TPC)

Material/Plant	Part used	Extract	IC <sub>50</sub> (mg/ml)	TPC $\pm$ SD
Acarbose	-	-	0.0009	NT
<i>Gentiana olivieri</i>	Aerial part	HA	0.1982	57.4 $\pm$ 2.7
<i>Helichrysum graveolens</i>	Capitulum	HA	0.7129	143.4 $\pm$ 9.4
		Aq	2.1979	92.9 $\pm$ 2.0
<i>H. plicatum</i> ssp. <i>plicatum</i>	Capitulum	HA	0.8570	139.5 $\pm$ 6.5
		Aq	5.0933	85.6 $\pm$ 15.7
<i>Juniperus communis</i> var. <i>saxatilis</i>	Fruit	HA	0.0044	21.0 $\pm$ 10.1
	Leaf	HA	0.0843	212.1 $\pm$ 9.9
		HA	-	4.8 $\pm$ 2.2
<i>J. oxycedrus</i> ssp. <i>oxycedrus</i>	Fruit	Aq	0.8054	24.8 $\pm$ 0.7
		HA	0.0473	191.0 $\pm$ 1.3
	Leaf	Aq	0.2606	160.4 $\pm$ 2.7
<i>Viscum album</i> ssp. <i>album</i>	Aerial part	HA	0.7962	21.2 $\pm$ 2.0
		Aq	3.7411	32.0 $\pm$ 0.2
<i>V. album</i> ssp. <i>austriacum</i>	Aerial part	HA	0.6653	35.8 $\pm$ 1.3
		Aq	1.3583	47.9 $\pm$ 0.8

Total phenol content data is expressed in mg equivalent of gallic acid (GAE) to 1 g of extract  
SD: Standard deviation, NT: Not tested, -:no activity, HA: Hydro-alcoholic, Aq: Aqueous

### Statistical analysis

All analyses were carried out in triplicates and the results were averaged. All values are expressed as the mean  $\pm$  standard deviation (SD); linear regression analyses and IC<sub>50</sub> calculations were done by using SigmaPlot 12.0 software. Microsoft Excel software was used to calculate correlation coefficients to determine the relationship between 2 variables.

## Results

### $\alpha$ -amylase inhibitory activity

$\alpha$ -Amylase inhibitory activities of the plant extracts were evaluated at 4 different logarithmic doses (3000, 1000, 300, 100  $\mu$ g/ml) and results were given in Table 2. All extracts except *H. plicatum* aqueous extract, showed a dose dependent inhibitory effect on  $\alpha$ -amylase enzyme. All the extracts exerted inhibitory activity at tested doses in varying proportions (3.5 – 55.7 % at 3000  $\mu$ g/ml). *H. graveolens* hydro-alcoholic extract exhibited the highest inhibitory activity at 3000  $\mu$ g/ml (55.7 %), while the inhibition percentage of the reference drug Acarbose was found to be 73.7 %. On the other hand, *J. oxycedrus* ssp. *oxycedrus* leaf hydroalcoholic extract possessed a continuous inhibitory effect on  $\alpha$ -amylase enzyme between 100–3000  $\mu$ g/ml (25.0–51.7%).

### $\alpha$ -glucosidase inhibitory activity

$\alpha$ -Glucosidase inhibitory activities of the plant extracts were evaluated at 5 different logarithmic doses between 0.3–10000  $\mu$ g/ml; the calculated IC<sub>50</sub> values are given in Table 3. All the extracts showed dose dependent inhibitory effect on  $\alpha$ -glucosidase enzyme. *J. communis* var. *saxatilis* fruit hydro-alcoholic extract possessed the highest inhibitory effect and its IC<sub>50</sub> value was found to be the lowest (IC<sub>50</sub> = 0.0044 mg/ml) among all extracts. IC<sub>50</sub> value of reference drug (Acarbose) was 0.0009 mg/ml. *H. plicatum* ssp. *plicatum* capitulum aqueous extract (IC<sub>50</sub> = 5.0933 mg/ml) and *V. album* ssp. *album* aerial

part aqueous extract (IC<sub>50</sub> = 3.7411 mg/ml) exerted the lowest enzyme inhibitory activity.

### ABTS radical cation scavenging activity

ABTS radical cation decolorization assay is a useful method for determining the antioxidant capacity of hydrogen donating antioxidants. ABTS<sup>+</sup> is a blue chromophore produced by the reaction between ABTS salt and potassium per-sulfate (26). After addition of extracts to ABTS radical cation, a strong reduction was observed and the blue color turned to white immediately in some extracts (*H. plicatum* capitulum hydro-alcoholic, *J. communis* var. *saxatilis* leaf hydro-alcoholic, *J. oxycedrus* ssp. *oxycedrus* leaf aqueous and hydro-alcoholic extracts) at 3000  $\mu$ g/ml concentration. Trolox used as a positive control, showed ABTS radical cation scavenging activity at all tested concentrations (100–3000  $\mu$ g/ml). *J. oxycedrus* ssp. *oxycedrus* fruit aqueous extract exerted the lowest radical scavenging activity (0–7.5%). The results of ABTS radical cation decolorization assay is given in Table 4.

### Total phenol content

Total phenol contents of all the extracts were measured and the results were shown in Table 3. The highest total phenol content was found in *J. communis* var. *saxatilis* leaf hydro-alcoholic extract (212.1  $\pm$  9.9 mg GAE/1 g extract) while the lowest was found in *J. oxycedrus* ssp. *oxycedrus* fruit hydro-alcoholic extract (4.8  $\pm$  2.2 mg GAE/1 g extract). Results presented in Table 3 show that there is a positive correlation between total phenol contents and ABTS radical scavenging activity of plant extracts (correlation coefficient=  $r = 0.8875$  at 3000  $\mu$ g/ml). However, the extracts with potent antioxidant activity and rich in phenolics did not show high inhibition on digestion enzymes. No correlation was observed between total phenol content and  $\alpha$ -amylase/ $\alpha$ -glucosidase inhibitory activity ( $r = 0.3959$  and  $r = 0.1669$  at 3000  $\mu$ g/ml respectively). Additionally correlation between radical

**Table 4.** ABTS radical scavenging activities of plant extracts

Material/Plant	Part used	Extract	Inh. % $\pm$ SD ( $\mu\text{g/ml}$ )			
			3000	1000	300	100
Trolox	-	-	>100	99.5 $\pm$ 0.25	38.9 $\pm$ 1.12	7.5 $\pm$ 0.72
<i>Gentiana olivieri</i>	Aerial part	HA	54.3 $\pm$ 1.20	20.5 $\pm$ 2.41	7.6 $\pm$ 1.03	2.6 $\pm$ 0.29
<i>Helichrysum graveolens</i>	Capitulum	Aq	77.0 $\pm$ 0.7	36.5 $\pm$ 0.74	11.2 $\pm$ 0.7	4.3 $\pm$ 0.91
		HA	88.5 $\pm$ 1.89	42.6 $\pm$ 1.95	15.0 $\pm$ 0.40	6.5 $\pm$ 2.53
<i>H. plicatum</i>	Capitulum	Aq	75.7 $\pm$ 0.61	36.1 $\pm$ 0.64	16.3 $\pm$ 0.8	2.7 $\pm$ 0.36
		HA	98.4 $\pm$ 2.66	53.5 $\pm$ 0.67	19.5 $\pm$ 0.36	6.8 $\pm$ 0.79
<i>Juniperus communis</i> var. <i>saxatilis</i>	Leaf	HA	99.5 $\pm$ 0.35	68.9 $\pm$ 1.03	24.3 $\pm$ 6.37	-
	Fruit	HA	42.5 $\pm$ 1.2	12.2 $\pm$ 0.57	-	-
<i>J. oxycedrus</i> ssp. <i>oxycedrus</i>	Fruit	Aq	7.5 $\pm$ 0.96	-	-	-
		HA	48.9 $\pm$ 0.55	19.4 $\pm$ 1.81	5.1 $\pm$ 0.70	2.3 $\pm$ 2.77
	Leaf	Aq	97.8 $\pm$ 0.83	37.2 $\pm$ 0.89	5.6 $\pm$ 0.5	-
<i>Viscum album</i> ssp. <i>album</i>	Aerial part	HA	97.8 $\pm$ 0.25	46.0 $\pm$ 0.51	4.7 $\pm$ 1.40	-
		Aq	33.7 $\pm$ 1.19	12.9 $\pm$ 2.05	-	-
		HA	50.2 $\pm$ 1.64	21.7 $\pm$ 2.78	13.3 $\pm$ 0.7	5.7 $\pm$ 1.44
<i>V. album</i> ssp. <i>austriacum</i>	Aerial part	Aq	47.5 $\pm$ 1.45	7.03 $\pm$ 0.06	-	-
		HA	56.6 $\pm$ 0.45	16.7 $\pm$ 0.7	8.9 $\pm$ 1.1	-

n=3, SD: standard deviation, -: no activity, HA:Hydro-alcoholic, Aq:Aqueous

scavenging and enzyme inhibitory activities of tested plant extracts were examined. No correlation was observed between ABTS radical scavenging and  $\alpha$ -amylase/ $\alpha$ -glucosidase inhibitory activity ( $r = -0.0876$  at 3000  $\mu\text{g/ml}$  and  $r = -0.1175$  respectively).

## Discussion

Hyperglycemia has been a classical risk factor in the development of diabetes and its complications. Therefore, control of blood glucose levels is critical in the early treatment of diabetes mellitus. One of the important therapeutic approaches is the prevention of carbohydrate absorption after food intake, which is facilitated by inhibition of the enteric enzymes including  $\alpha$ -glucosidase and  $\alpha$ -amylase present in the intestinal brush border (27, 28). The inhibition of these enzymes has been a strong option in the prevention of diabetes. So, inhibitors like Acarbose, voglibose, and miglitol are widely used in type 2 diabetic patients nowadays. Moreover, studies are being carried out to find new amylase and glucosidase inhibitors from natural sources (29–31).

The aim of this study is to clarify the mechanism of action of selected plants on carbohydrate metabolism. For this purpose, the inhibitory effect of 15 extracts obtained from different parts of 7 plants on  $\alpha$ -glucosidase and  $\alpha$ -amylase were assessed and compared with the  $\alpha$ -glucosidase inhibitor, Acarbose. Also, radical scavenging activity and total phenol content of the extracts were investigated.

In our previous studies, we demonstrated significant hypoglycaemic and antidiabetic activities of hydro-alcoholic extracts of *H. graveolens* capitulum, *J. oxycedrus* ssp. *oxycedrus* leaves and *J. communis* var. *saxatilis* fruits in normoglycaemic, glucose loaded and streptozotocin-induced diabetic rats. In the present study, these extracts which were found to have potent antidiabetic activity, have also shown high inhibitory effect on enzymes that have an important role in carbohydrate metabolism. There was no correlation between total phenol

content and  $\alpha$ -amylase/ $\alpha$ -glucosidase inhibitory activity of these plant extracts.

Many studies were conducted on the chemical profile of the selected medicinal plants. Orhan *et al* (13, 14) isolated and identified many compounds that are responsible for the antidiabetic activity of *J. oxycedrus* ssp. *oxycedrus* (Joso). Through *in vivo* bioactivity-guided fractionation processes, shikimic acid, 4-O- $\beta$ -D-glucopyranosyl ferulic acid, and oleuropeic acid-8-O- $\beta$ -D-glucopyranoside were isolated from the active subfractions of Joso fruit hydro-alcoholic extracts as the active components (14). Jeong *et al* (2012) showed strong inhibitory effects of ferulic acid derivatives on  $\alpha$ -amylase and  $\alpha$ -glucosidase enzymes (32). Therefore, we propound that  $\alpha$ -amylase inhibitory effect of Joso fruit hydro-alcoholic extract might be produced by the presence of ferulic acid and other chemical constituents. Additionally, the major antidiabetic compounds in subfractions of Joso leaves were identified as fatty acids such as palmitic, linoleic, and linolenic acid (13). Su *et al* (2013) investigated the inhibitory mechanisms of fatty acids on key enzymes related to type 2 diabetes. Oleic and linoleic acids were found to have potent inhibitory effects on  $\alpha$ -glucosidase activity (33). Thus, fatty acids might contribute to the  $\alpha$ -glucosidase enzyme inhibitory effects of other active compounds found in the Joso leaf hydro-alcoholic extract.

Leaves and fruits of *J. communis* var. *saxatilis* contain relatively high amounts of monoterpene hydrocarbons such as  $\alpha$ -pinene, limonene and  $\beta$ -myrcene (34). The main monoterpene component for these parts of the plant was  $\alpha$ -pinene. Bařak and Candan (2013) found that  $\alpha$ -pinene in *Laurus nobilis* essential oil inhibited  $\alpha$ -glucosidase (35). On the other hand, *J. communis* var. *saxatilis* leaf hydroalcoholic extract showed significantly  $\alpha$ -amylase inhibitory effect which may be due to the presence of some secondary metabolites such as lignans, coumarins, sterols, aliphatic compounds, and other terpenes in the hydroalcoholic extract (34).

Results of  $\alpha$ -amylase inhibitory activity assay showed that *H. graveolens* hydro-alcoholic extract has *in vitro* enzyme inhibition in a degree similar to Acarbose at 3000  $\mu\text{g/ml}$ . Flavonoids, acetophenones, phloroglucinol, pyrones, triterpenoids, and sesquiterpenes are secondary metabolites of the genus *Helichrysum* (36). Additionally, Albayrak *et al* (2010) reported the presence of chlorogenic acid, caffeic acid, ferulic acid, syringic acid, apigenin, apigenin-7-glucoside, and hesperidin; luteolin, naringenin, quercetin, resveratrol in the methanol extracts of *H. graveolens*, and chlorogenic acid were found to be the major phenolics in the extract (37). Narita *et al* (2008) reported the strong inhibitory effect of chlorogenic acid and its derivatives on porcine pancreas  $\alpha$ -amylase (38). It is considered that high phenolic content (143.4 mg GAE/1 g extract) of hydro-alcoholic extract of *H. graveolens* capitulum might support the enzyme inhibitory effect of other constituents like chlorogenic acid and its derivatives.

*H. plicatum* capitulum hydro-alcoholic, *J. communis* var. *saxatilis* leaf hydro-alcoholic, and *J. oxycedrus* ssp. *oxycedrus* leaf aqueous and hydro-alcoholic extracts have shown strong ABTS radical cation scavenging activity. Antioxidant effects of these plants might cooperate with their antidiabetic activity and these plants might be a better choice for complementary remedies for type 2 diabetic patients.

## Conclusion

This is the first study on the *in vitro* antidiabetic activities of these seven plants: *G. olivieri*, *H. graveolens*, *H. plicatum* ssp. *plicatum*, *J. oxycedrus* ssp. *oxycedrus*, *J. communis* var. *saxatilis*, and *V. album* (ssp. *album* and ssp. *austriacum*). These seven plants with previously reported *in vivo* antidiabetic effect were tested for enzyme inhibitory and radical scavenging activities. Among these, *H. graveolens* hydro-alcoholic extract, *J. communis* leaf hydro-alcoholic extract and *J. oxycedrus* leaf and fruit hydro-alcoholic extracts were found to have inhibitory effect on  $\alpha$ -amylase. On the other hand, hydroalcoholic extracts of *J. communis* (leaf, fruit) *J. oxycedrus* (leaf) had potent inhibitory activity on  $\alpha$ -glucosidase. In conclusion, the findings of this investigation indicate that these plants might be ameliorate hyperglycemia in type 2 diabetics by their inhibitory effect on  $\alpha$ -glucosidase and  $\alpha$ -amylase. It is concluded that further studies are needed to explain the mechanism of actions of the other extracts and their active constituents.

## Acknowledgment

This study was financially supported by the Research Fund of Gazi University (02/2011-22). We are thankful to Bayer Group for providing us with Acarbose.

## References

- Marles RJ, Farnsworth NR. Antidiabetic plants and their active constituents. *Phytomedicine* 1995; 2:137-189.
- Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010; 87:4-14.
- Fujisawa T, Ikegami H, Inoue K, Kawabata Y, Ogihara T. Effect of two  $\alpha$ -glucosidase inhibitors, voglibose and acarbose, on postprandial hyperglycemia correlates with subjective abdominal symptoms. *Metabolism* 2005; 54:387-390.
- Gray AM, Flatt PR. Nature's own pharmacy: the diabetes perspective. *Proc Nutr Soc* 1997; 56:507-517.
- Gholamhosseinian A, Falah H, Sharififar F, Mirtajaddini M. The inhibitory effect of some Iranian plants extracts on the alpha glucosidase. *Iran J Basic Med Sci* 2008; 11:1-9.
- World Health Organization. WHO Traditional Medicine Strategy 2002-2005, WHO, Geneva, Switzerland: 2002.
- Andrade-Cetto A, Heinrich M. Mexican plants with hypoglycaemic effect used in the treatment of diabetes. *J Ethnopharmacol* 2005; 99:325-348.
- Katemo M, Mpiana PT, Mbala BM, Mihigo SO, Ngbolua KN, Tshibangu DS, *et al*. Ethnopharmacological survey of plants used against diabetes in Kisangani City (DR Congo). *J Ethnopharmacol* 2012; 144:39-43.
- Tahraoui A, El-Hilaly J, Israili ZH, Lyoussi B. Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in south-eastern Morocco (Errachidia province). *J Ethnopharmacol* 2007; 110:105-117.
- Yaniv Z, Dafni A, Friedman J, Palevitch D. Plants used for the treatment of diabetes in Israel. *J Ethnopharmacol* 1987; 19:145-151.
- Sezik E, Aslan M, Yeşilada E, Ito S. Hypoglycaemic activity of *Gentiana olivieri* and isolation of the active constituent through bioassay-directed fractionation techniques. *Life Sci* 2005; 76:1223-1238.
- Aslan M, Deliorman Orhan D, Orhan N, Sezik E, Yeşilada E. A study of antidiabetic and antioxidant effects of *Helichrysum graveolens* capitulum in streptozotocin-induced diabetic rats. *J Med Food* 2007; 10:396-400.
- Orhan N, Aslan M, Demirci B, Ergun F. A bioactivity guided study on the antidiabetic activity of *Juniperus oxycedrus* subsp. *oxycedrus* L. leaves. *J Ethnopharmacol* 2012; 140:409-415.
- Orhan N, Aslan M, Pekcan M, Deliorman Orhan D, Bedir E, Ergun F. Identification of hypoglycaemic compounds from berries of *Juniperus oxycedrus* subsp. *oxycedrus* through bioactivity guided isolation technique. *J Ethnopharmacol* 2012; 139:110-118.
- Deliorman Orhan D, Aslan M, Şendoğdu N, Ergun F, Yeşilada E. Evaluation of the hypoglycemic effect and antioxidant activity of three *Viscum album* subspecies (European mistletoe) in streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2005; 98:95-102.
- Aslan M, Deliorman Orhan D, Orhan N, Sezik E, Yeşilada E. *In vivo* antidiabetic and antioxidant potential of *Helichrysum plicatum* ssp. *plicatum* capitulum in streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2007; 109:54-59.

17. Hudecová A, Kusznierevicz B, Hašplová K, Huk A, Magdolenová Z, Miadoková E, et al. *Gentiana asclepiadea* exerts antioxidant activity and enhances DNA repair of hydrogen peroxide- and silver nanoparticles-induced DNA damage. *Food Chem Toxicol* 2012; 50:3352-3359.
18. Loizzo MR, Tundis R, Conforti F, Saab AM, Statti GA, Menichini F. Comparative chemical composition, antioxidant and hypoglycaemic activities of *Juniperus oxycedrus* ssp. *oxycedrus* L. berry and wood oils from Lebanon. *Food Chem* 2007; 105:572-578.
19. Kizilarıslan Ç, Özhatay N. An ethnobotanical study of the useful and edible plants of İzmit. *Marmara Pharm J* 2012; 16:194-200.
20. Maritim AC, Sanders RA, Watkins JB. Diabetes, oxidative stress, and antioxidants: A review. *J Biochem Mol Toxicol* 2003; 17:24-38.
21. Ali H, Houghton PJ, Soumyanath A.  $\alpha$ -Amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*. *J Ethnopharmacol* 2006; 107:449-455.
22. Lam SH, Chen JM, Kang CJ, Chen CH, Lee SS.  $\alpha$ -Glucosidase inhibitors from the seeds of *Syagrus romanzoffiana*. *Phytochemistry* 2008; 69:1173-1178.
23. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999; 26:1231-1237.
24. Meot-Duros L, Le Floch G, Magne C. Radical scavenging, antioxidant and antimicrobial activities of halophytic species. *J Ethnopharmacol* 2008; 116:258-262.
25. Gao X, Ohlander M, Jeppsson N, Björk L, Trajkovski V. Changes in antioxidant effects and their relationship to phytonutrients in fruits of sea Buckthorn (*Hippophae rhamnoides* L.) during maturation. *J Agric Food Chem* 2000; 48:1485-1490.
26. Mayur B, Sandesh S, Shruti S, Sung-Yum S. Antioxidant and  $\alpha$ -glucosidase inhibitory properties of *Carpesium abrotanoides* L. *J Med Plant Res* 2010; 4:1547-1553.
27. Inzucchi SE. Oral antihyperglycemic therapy for type 2 diabetes: scientific review. *J Am Med Assoc* 2002; 287:360-372.
28. Toeller M. Alpha-glucosidase inhibitors in diabetes: efficacy in NIDDM subjects. *Eur J Clin Invest* 1994; 24:31-35.
29. Bhat M, Zinjarde SS, Bhargava SY, Kumar AR, Joshi BN. Antidiabetic Indian plants: a good source of potent amylase inhibitors. *Evid-Based Compl Alt Med* 2011; 2011:810207.
30. Kumarappan C, Mandal SC.  $\alpha$ -Glucosidase inhibitory activity and in vitro antioxidant activities of alcohol-water extract (AWE) of *Ichnocarpus frutescens* leaves. *Med Chem Res* 2008; 17:219-233.
31. Matsui T, Ebuchi S, Kobayashi M, Fukui K, Sugita K, Terahara N. Anti-hyperglycemic effect of diacylated anthocyanin derived from *Ipomea batatas* cultivar Ayamurasaki can be achieved by through the  $\alpha$ -glucosidase inhibitory action. *J Agric Food Chem* 2002; 50:7244-7248.
32. Jeong EY, Cho KS, Lee HS.  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitors isolated from *Triticum aestivum* L. sprouts. *J Korean Soc App Biol Chem* 2012; 55:47-51.
33. Su CH, Hsu CH, Ng LT. Inhibitory potential of fatty acids on key enzymes related to type 2 diabetes. *Biofactors* 2013; 39:415-421.
34. Lohani H, Haider SZ, Chauhan NK, Sah S, Andola HC. Aroma profile of two *Juniperus* species from Alpine region in Uttarakhand. *J Nat Prod* 2013; 6:38-43.
35. Basak SS, Candan F. Effect of *Laurus nobilis* L. essential oil and its main components on  $\alpha$ -glucosidase and reactive oxygen species scavenging activity. *Iran J Pharm Res* 2013; 12:367-379.
36. Rosa A, Deiana M, Atzeri A, Corona G, Incani A, Melis MP, et al. Evaluation of the antioxidant and cytotoxic activity of arzanol, a prenylated  $\alpha$ -pyrone-phloroglucinol etherodimer from *Helichrysum italicum* subsp. *microphyllum*. *Chem-Biol Interact* 2007; 165:117-126.
37. Albayrak S, Aksoy A, Sagdic O, Hamzaoglu E. Compositions, antioxidant and antimicrobial activities of *Helichrysum* (Asteraceae) species collected from Turkey. *Food Chem* 2010; 119:114-122.
38. Narita Y, Kimura R, Nakagiri O, Inouye K. Kinetic analysis and mechanism on the inhibition of chlorogenic acids against porcine pancreas  $\alpha$ -amylase. *Proceedings of the 22<sup>nd</sup> International Conference on Coffee Science (ASIC), 2008, Campinas, SP-Brazil* p. 171-175.