

Synthesis and Purification of 7-Prenyloxycoumarins and Herniarin as Bioactive Natural Coumarins

¹Mahdi Askari, ¹Amirhossein Sahebkar, *¹Mehrdad Iranshahi

Abstract

Objective(s)

7-prenyloxycoumarins including 7-isopentenylloxycoumarin, auraptene and umbelliprenin, and herniarin have been widely recognized as bioactive coumarins. This paper presents the ways to synthesis these compounds.

Materials and Methods

7-prenyloxycoumarins were synthesized by reaction between 7-hydroxycoumarin (1 M) and relevant prenyl bromides (1.5 M) in acetone at room temperature. The reaction was carried out in the presence of DBU (1, 8-diazabicyclo [5.4.0] undec-7-ene) (2 M). After 24 hr, the mixture was concentrated under reduced pressure. The compounds were purified by column chromatography.

Results

Three bioactive 7-prenyloxycoumarins, namely, umbelliprenin, auraptene and 7-isopentenylloxycoumarin, together with herniarin were synthesized from 7-hydroxycoumarin under alkaline conditions (DBU) and then purified by column chromatography. The structures of the products were characterized by NMR spectroscopic method including ¹H- and ¹³C-NMR experiments.

Conclusion

The method of synthesis for 7-prenyloxycoumarins and herniarin which is presented here has not been reported yet. Moreover, for the first time, umbelliprenin was chemically prepared in this work.

Keywords: Auraptene, Herniarin, 7- Isopentenylloxycoumarin, 7-Prenyloxycoumarins, Synthesis, Umbelliprenin

1- Department of Pharmacognosy, Biotechnology Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

*Corresponding author: Tel: +98-511-8823255-66; Fax: +98-511-8823251; email: Iranshahim@mums.ac.ir

Introduction

7-Prenyloxycoumarins are a group of secondary metabolites which are found mainly in plants belonging to the families of Rutaceae and Umbelliferae. These compounds possess various biological and pharmacological activities which have gained attention of researchers in the last two decades. Three of the most studied prenylated coumarins include auraptene, umbelliprenin and 7-isopentenylloxycoumarin.

Umbelliprenin (Figure 1) is a naturally occurring prenylated coumarin which is synthesized by various *Ferula* species (1-4). It has also been found in various plant species consumed as food or used for food preparation such as in celery, *Angelica archangelica*, *Coriandrum sativum* and *Citrus limon* (5). It has been reported that umbelliprenin inhibits the red pigment production in *Serratia marcescens* (2), inhibits squalene-hopene cyclase (SHC) (an enzyme taking part in sterol synthesis) (6), decreases matrix metalloproteinase (MMP) activity (4), exhibits antileishmanial activity against promastigotes (3), induces apoptosis in human M4Beu metastatic pigmented melanoma cells (7) and exerts cancer chemopreventive activity (8). Auraptene (7-geranyloxycoumarin) is the most abundant geranyloxycoumarin extracted from plants belonging to the genus *Citrus* (1), and is known as a potent cancer chemopreventive (8-10) and anti-tumor agent against many types of cancers (9-13). Moreover, it exerts anti-inflammatory activity (14, 15) and is capable of suppressing the release of tumor necrosis factor alpha (TNF- α) (16), superoxide anion generation by inflammatory leukocytes (17) and I κ B (nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor) degradation (18). Auraptene also causes complete inhibition of platelet aggregation, induced by arachidonic acid and platelet activated factor *in vitro* (19). Finally, auraptene and its analogs have been shown to exhibit spasmolytic activity (20, 21). 7-isopentenylloxycoumarin extracted from plants belonging to genus *Ruta* (1), has been shown to possess promising antifungal activity (22, 23). Besides, this compound exerts antitumor (23) and chemopreventive

activity (24). Recently, auraptene and 7-isopentenylloxycoumarin have been found to possess significant neuroprotective activity (25). Herniarin occurs commonly in higher plants, e.g. *Matricaria* spp. (26, 27) and *Lavandula* spp. (28). In a previous report, herniarin was introduced to have anti-dermatophytic activity (29).

Angioni and colleagues (30) have reported the preparation of auraptene with a reaction between 7-hydroxycoumarin and geranyl bromide in K₂CO₃ solution. In another work, auraptene was synthesized from umbelliferone by prenylation with NaH and geranyl bromide in DMF (31). Herniarin was also prepared via a reaction between ortho-methoxy phenol and alkynoates in the presence of a palladium catalyst (32).

In the present study, we wish to report a new synthesis of four bioactive coumarins, namely, umbelliprenin, auraptene, 7-isopentenylloxycoumarin and herniarin, and their characterization, using NMR spectroscopic methods including ¹H- and ¹³C-NMR spectroscopy.




Figure 1. Chemical structures of 7-prenyloxycoumarins: umbelliprenin (1), auraptene (2) and 7-isopentenylloxycoumarin (3), and herniarin (4).

Materials and Methods

Chemistry

Melting points were determined on a Sanyo Gallenkamp apparatus. NMR spectra were measured on a Bruker DRX AV 500 (Bruker Biospin, Rheinstetten, Germany). ¹H- and ¹³C-NMR spectrum were measured, using an inverse-detection probe (5 mm). The operating frequencies were 500.13 MHz for acquiring ¹H-NMR and 125.75 MHz for ¹³C-NMR spectra. Samples were measured at 300 K in CDCl₃ with TMS as the internal standard. Column chromatography was conducted with silica gel 230-400 mesh (Merck).

Preparation of umbelliprenin [(E, E)-7-(3, 7, 11-trimethyl-2, 6, 10-dodecatrienyloxy) chromen-2-one]

Umbelliprenin (7-farnesyloxycoumarin) was synthesized in 71% yield by reaction between 7-hydroxycoumarin (1 M) and *trans-trans-*

farnesyl bromide (1.5 M) in acetone at room temperature. The reaction was carried out in the presence of DBU (1, 8-diazabicyclo [5.4.0] undec-7-ene) (2 M). After 24 hr, the mixture was concentrated under reduced pressure. Umbelliprenin was purified by column chromatography (petroleum ether/ethyl acetate 9:1 v/v) as white crystals (mp = 57.5-59.1 °C) (5). The chemical structure of umbelliprenin is shown in Figure 1, and ¹H- and ¹³C-NMR data for this compound are presented in Tables 1 and 2.

Preparation of auraptene [7-((E)-3, 7-Dimethylocta-2, 6-dienyloxy)-2H-chromen-2-one]

Auraptene (7-geranyloxycoumarin) was similarly synthesized in 65% yield by reaction between 7-hydroxycoumarin (1 M) and *trans-geranyl* bromide (1.5 M) in acetone at room temperature. The reaction was carried out in the presence of DBU (2 M). After 24 hr, the

Table 1. ¹H-NMR data for compounds 1-4 (CDCl₃, 500 MHz)^a.

H	1	2	3	4
2	-	-	-	-
3	6.25 d (9.6)	6.24 d (9.6)	6.24 d (9.6)	6.25 d (9.6)
4	7.63 d (9.6)	7.63 d (9.6)	7.63 d (9.6)	7.62 d (9.6)
5	7.36 d (7.2)	7.36 d (7.2)	7.36 d (7.2)	7.37 d (7.2)
6	6.85 dd (7.2, 2.0)	6.85 dd (7.2, 2.0)	6.85 dd (7.2, 2.0)	6.85 dd (7.2, 2.0)
7	-	-	-	-
8	6.82 d (2.0)	6.82 d (2.0)	6.82 d (2.0)	6.82 d (2.0)
9	-	-	-	-
10	-	-	-	-
1'	4.60 d (7.0)	4.59 d (7.0)	4.57 d (7.0)	3.86 d (7.0)
2'	5.47 t (7.0)	5.47 t (7.0)	5.47 t (7.0)	-
3'	-	-	-	-
4'	2.12 m	2.13 m	1.80 m	-
5'	2.15 m	2.15 m	1.76 m	-
6'	5.07 q (7.0)	5.08 q (7.0)	-	-
7'	-	-	-	-
8'	1.97 m	1.75 m	-	-
9'	2.05 m	1.65 m	-	-
10'	5.09 q (7.0)	1.59 q (7.0)	-	-
11'	-	-	-	-
12'	1.76 s	-	-	-
13'	1.60 s	-	-	-
14'	1.67 s	-	-	-
15'	1.59 s	-	-	-

^aJ values are in parenthesis and reported in Hz; chemical shifts are given in ppm.

mixture was concentrated under reduced pressure. Auraptene was easily purified by column chromatography (petroleum ether/ethyl acetate 9:1 v/v) as white crystals (mp = 62.7-63.4 °C) (5).

The chemical structure of auraptene is shown in Figure 1, and ¹H- and ¹³C-NMR data for this compound are presented in Tables 1 and 2.

Table 2. ¹³C-NMR data for compounds 1-4 (CDCl₃, 125.7 MHz).

C	1	2	3	4
2	161.2	161.2	161.2	161.1
3	112.9	112.9	112.9	112.5
4	143.4	143.4	143.4	143.3
5	128.6	128.6	128.6	128.7
6	113.2	113.2	113.1	113.0
7	162.1	162.1	162.0	162.8
8	101.5	101.5	101.5	100.8
9	155.8	155.8	155.8	155.8
10	112.4	112.3	112.3	112.5
1'	65.4	65.4	65.3	55.7
2'	118.4	118.3	118.5	-
3'	142.3	142.3	139.2	-
4'	39.5	39.4	25.7	-
5'	26.1	26.4	18.2	-
6'	123.4	123.5	-	-
7'	135.5	131.9	-	-
8'	39.6	16.7	-	-
9'	26.6	25.6	-	-
10'	124.2	17.6	-	-
11'	131.3	-	-	-
12'	16.7	-	-	-
13'	16	-	-	-
14'	25.6	-	-	-
15'	17.6	-	-	-

Preparation of 7-isopentenylcoumarin [7-(isopentenyl) chromen-2-one]

7-isopentenylcoumarin was also synthesized in 45% yield by reaction between 7-hydroxycoumarin (1 M) and isopentenyl bromide (1.5 M) in acetone at room temperature. The reaction was carried out in the presence of DBU (2 M). After 24 hr, the mixture was concentrated under reduced pressure. 7-isopentenylcoumarin was purified by column chromatography (petroleum ether/ethyl acetate 9:1 v/v) as white crystals (mp = 74.3-75.2 °C) (5). The chemical structure of 7-isopentenylcoumarin is shown in Figure 1, and ¹H- and ¹³C-NMR data for this compound are presented in Tables 1 and 2.

Preparation of herniarin (7-methoxychromen-2-one)

Herniarin (7-methoxycoumarin) was synthesized in 54.8% yield by reaction between 7-hydroxycoumarin (1 M) and methyl

iodide (1.5 M) in acetone at room temperature. The reaction was carried out in the presence of DBU (2 M). After 24 hr, the mixture was concentrated under reduced pressure. Herniarin was purified by column chromatography (petroleum ether/ethyl acetate 9:1 v/v) as white crystals (mp= 120.0-121.0 °C) (5). The chemical structure of herniarin is shown in Figure 1, and ¹H- and ¹³C-NMR data for this compound are presented in Tables 1 and 2.

Results and Discussion

To our knowledge, the method of synthesis presented in the current study has not been reported to date, especially for umbelliprenin that has not been previously chemically prepared. However, Angioni and colleagues (30) reported the preparation of auraptene with a reaction between 7-hydroxycoumarin and geranyl bromide in K₂CO₃ solution (acetone). The yield of the mentioned reaction was 85% after 10 hr. In another work, auraptene was

synthesized from umbelliferone by prenylation with NaH and geranyl bromide in DMF (31). Trost and colleagues (32) recently reported a new method for the synthesis of herniarin and other coumarins in the presence of palladium catalyst. In our work, we studied the prenylation of 7-hydroxycoumarin with different prenyl moieties (isopentenyl, geranyl and farnesyl) by, using DBU as an alkaline reagent in acetone. The yield of each reaction is reported in the experimental section. Besides, the synthesis of herniarin as a non-prenylated coumarin has also been reported in this paper.

The ^1H - and ^{13}C -NMR data, together with melting points of the four synthesized compounds namely auraptene, umbelliprenin, 7-isopentenylloxycoumarin and herniarin were in agreement with those previously described in the literature (5, 33, 34).

The ^1H -NMR spectrum of umbelliprenin showed resonances characteristic for four methyl singlets at δ_{H} 1.59 (H-15'), 1.60 (H-13'), 1.67 (H-14') and 1.76 (H-12'), and eight methine resonances at δ_{H} 5.07 (H-6', t, $J = 7.0$ Hz), 5.09 (H-10', t, $J = 7.0$ Hz), 5.47 (H-2', t, $J = 7.0$ Hz), 6.25 (H-3, d, $J = 9.6$ Hz), 6.82 (H-8, d, $J = 2.0$ Hz), 6.85 (H-6, dd, $J = 7.2$ Hz, $J = 2$ Hz), 7.36 (H-5, d, $J = 7.2$ Hz) and 7.63 (H-4, d, $J = 9.6$ Hz). Three aromatic protons at δ_{H} 6.82 (H-8), 6.85 (H-6) and 7.36 (H-5) suggested the presence of a 7, 9, 10-trisubstituted benzene ring, which was supported by the ^{13}C -NMR spectrum.

The ^{13}C -NMR resonances displayed 24 carbon signals, nine being typical of an umbelliferone skeleton and the other 15 signals were ascribable to a prenyl (farnesyl) moiety. The ^{13}C -NMR spectrum showed four methyl signals at δ_{C} 16.0 (C-13'), 16.7 (C-12'), 17.6 (C-15') and 25.6 (C-14'), three methylene

signals at δ_{C} 26.1 (C-5'), 26.6 (C-9') and 65.4 (C-1'), eight methine signals at δ_{C} 101.5 (C-8), 112.9 (C-3), 118.4 (C-2'), 113.2 (C-6), 124.2 (C-10'), 123.4 (C-6'), 128.6 (C-5) and 143.4 (C-4), and seven quaternary carbon signals at δ_{C} 112.4 (C-10), 131.3 (C-11'), 135.5 (C-7'), 155.8 (C-9), 142.3 (C-3'), 161.2 (C-2) and 162.1 (C-7), including one carbonyl function which was indicated by the downfield signal at δ_{C} 161.2 (C-2). These ^1H - and ^{13}C -NMR data were in accordance with those previously reported for umbelliprenin (5, 33).

The ^1H - and ^{13}C -NMR spectra of other coumarins including auraptene, 7-isopentenylloxycoumarin and herniarin were in agreement with the literature (34). Their structures were similarly characterized. The full NMR data of synthesized coumarins are shown in Table 1.

Conclusion

7-prenyloxycoumarins are easily synthesized via a reaction between prenyl bromides and umbelliferone in an alkaline condition (DBU in acetone) and then are successfully purified with silica column chromatography. Although, no by products were detected in the reaction, the reaction was not accomplished after 48 hr. It might be due to impurities of the reagents or other factors.

Finally, it is worth noting that the method of synthesis for 7-prenyloxycoumarins and herniarin which is presented here has not been reported yet. Moreover, for the first time umbelliprenin was chemically prepared in the present work.

Acknowledgments

This research was supported by a grant (186336) from Mashhad University of Medical Sciences Research Council, Mashhad, Iran.

References

1. Curini M, Cravotto G, Epifano F, Giannone G. Chemistry and biological activity of natural and synthetic prenyloxycoumarins. *Curr Med Chem* 2006;13:199-222.
2. Iranshahi M, Shahverdi AR, Mirjani R, Amin G, Shafiee A. Umbelliprenin from *Ferula persica* roots inhibits the red pigment production in *Serratia marcescens*. *Z Naturforsch* 2004; 59c:506-508.
3. Iranshahi M, Arfa P, Ramezani M, Jaafari MR, Sadeghian H, Bassarello C, *et al.* Sesquiterpene coumarins from *Ferula szowitsiana* and *in vitro* antileishmanial activity of 7-prenyloxycoumarins against promastigotes. *Phytochemistry* 2007; 68:554-561.

4. Shahverdi AR, Saadat F, Khorramizadeh MR, Iranshahi M, Khoshayand MR. Two matrix metalloproteinases inhibitors from *Ferula persica* var. *persica*. *Phytomedicine* 2006; 13:712-717.
5. Murray RDH, Mendez J, Brown RA. *The Natural Coumarins*. New York: John Wiley & Sons; 1982.
6. Cravotto G, Balliano G, Robaldo B, Oliaro-Bosso S, Chimichi S, Boccalini M. Farnesyloxy coumarins, a new class of squalene-hopene cyclase inhibitors. *Bioorg Med Chem Lett* 2004; 14:1931-1934.
7. Barthomeuf C, Lim S, Iranshahi M, Chollet P. Umbelliprenin from *Ferula szowitsiana* inhibits the growth of human M4Beu metastatic pigmented melanoma cells through cell-cycle arrest in G1 and induction of caspase-dependent apoptosis. *Phytomedicine* 2008; 15:103-111.
8. Iranshahi M, Kalategi F, Rezaee R, Shahverdi AR, Ito C, Furukawa H, *et al*. Cancer chemopreventive activity of terpenoid coumarins from *Ferula* species. *Planta Med* 2008; 74:147-150.
9. Tanaka T, Kawabata K, Kakumoto M, Mastunaga K, Mori H, Murakami A, *et al*. Chemoprevention of 4-nitroquinoline 1-oxide-induced oral carcinogenesis by *Citrus aurapetne* in rats. *Carcinogenesis* 1998; 19:425-431.
10. Tanaka T, Kawabata K, Kakumoto M, Hara A, Murakami A, Kuki W, *et al*. *Citrus aurapetne* exerts dose-dependent chemopreventive activity in rat large bowel tumorigenesis: the inhibition correlates with suppression of cell proliferation and lipid peroxidation and with induction of phase II drug-metabolizing enzymes. *Cancer Res* 1998; 58:2550-2556.
11. Sakata K, Hara A, Hirose Y, Yamada Y. Dietary supplementation of the *Citrus* antioxidant auraptene inhibits N,N-Diethylnitrosamine-Induced rat hepatocarcinogenesis. *Oncology* 2004; 66:244-252.
12. Murakami A, Kuki W, Takahashi Y, Yonei H, Nakamura Y, Ohto Y, *et al*. Auraptene, a citrus coumarin, inhibits 12-O-tetradecanoylphorbol-13-acetate-induced tumor promotion in ICR mouse skin, possibly through suppression of superoxide generation in leukocytes. *Jpn J Cancer Res* 1997; 88:443-452.
13. Kawabata K, Tanaka T, Yamamoto T, Hara A, Murakami A, Koshimizu K, *et al*. Suppression of N-nitrosomethylbenzylamine-induced rat esophageal tumorigenesis by dietary feeding of auraptene. *J Exp Clin Cancer Res* 2000; 19:45-52.
14. Curini M, Epifano F, Maltese F, Marcotullio MC, Tubaro A, Altinier G, *et al*. Synthesis and anti-inflammatory activity of natural and semisynthetic geranyloxy coumarins. *Bioorg Med Chem Lett* 2004; 14:2241-2243.
15. Murakami A, Nakamura Y, Ohto Y, Yano M, Ohigashi H. Suppressive effects of citrus fruits on free radical generation and nobiletin, an anti-inflammatory polymethoxyflavonoid. *BioFactors* 2000; 12:187-192.
16. Tanaka T, Sugiura H, Inaba R, Nishikawa A, Murakami A, Koshimizu K, *et al*. Immunomodulatory action of citrus auraptene on macrophage functions and cytokine production of lymphocytes in female BALB/c mice. *Carcinogenesis* 1999; 20:1471-1476.
17. Murakami A, Nakamura Y, Tanaka T, Kawabata K, Takahashi K, Koshimizu K, *et al*. Suppression by citrus auraptene of phorbol ester- and endotoxin-induced inflammatory responses: role of attenuation of leukocyte activation. *Carcinogenesis* 2000; 21:1843-1850.
18. Murakami A, Matsumoto K, Koshimizu K, Ohigashi H. Effects of selected food factors with chemopreventive properties on combined lipopolysaccharide- and interferon-g-induced IKB degradation in RAW264.7 macrophages. *Cancer Lett* 2003; 195:17-25.
19. Chen IS, Lin YC, Tsai IL, Teng CM, Ko FN, Ishikawa T, *et al*. Coumarins and anti-platelet aggregation constituents from *Zanthoxylum schinifolium*. *Phytochemistry*. 1995; 39:1091-1097.
20. Yamada Y, Nakatani N, Fuwa H. Spasmolytic activity of geranyloxy coumarin-related compounds. *Agric Biol Chem* 1987; 51:1711-1713.
21. Yamada Y, Okamoto M, Kikuzaki H, Nakatani N. Spasmolytic activity of auraptene analogs. *Biosci Biotechnol Biochem* 1997; 61:740-742.
22. Hamerski D, Schmitt D, Matern U. Induction of two prenyltransferases for the accumulation of coumarin phytoalexins in elicitor-treated *Ammi majus* cell suspension cultures. *Phytochemistry* 1990; 29:1131-1135.
23. Rahalison L, Benathan M, Monod M, Frenk E, Gupta MP, Solis PN, *et al*. Antifungal principles of *Baccharis pedunculata*. *Planta Med* 1995; 61:360-362.
24. Baba M, Jin Y, Mizuno A, Suzuki H, Okada Y, Takasuka N, *et al*. Studies on cancer chemoprevention by traditional folk medicines XXIV. Inhibitory effect of a coumarin derivative, 7-isopentenylcoumarin, against tumor-promotion. *Biol Pharm Bull* 2002; 25:244-246.
25. Epifano F, Molinaro G, Genovese S, Ngomba RT, Nicoletti F, Curini M. Neuroprotective effect of prenyloxy coumarins from edible vegetables. *Neurosci Lett* 2008; 443:57-60.
26. Ahmad A, Misra LN. Isolation of herniarin and other constituents from *Matricaria chamomilla* flowers. *Int J Pharm* 1997; 35:121-125.
27. Ma CM, Winsor L, Daneshtalab M. Quantification of spiroether isomers and herniarin of different parts of *Matricaria matricarioides* and flowers of *Chamaemelum nobile*. *Phytochem Anal* 2007; 18:42-49.
28. Brown SA. Biosynthesis of the coumarins IV. The formation of coumarin and herniarin in *Lavander*. *Phytochemistry* 1963; 2:137-144.
29. Mares D, Romagnoli C, Bruni A. Antidermatophytic activity of herniarin in preparations of *Chamomilla recutita* (L.) Rauschert. *Plantas Med Phytother* 1993; 26:91-100.

Synthesis of 7-prenyloxycoumarins

30. Angioni A, Cabras G, D'hallewin G, Pirsì FM, Reniero F, Schirra M. Synthesis and inhibitory activity of 7-geranyloxycoumarin against *Penicillium* species in *Citrus* fruit. *Phytochemistry* 1998; 47:1521-1525.
31. Curini M, Epifano F, Maltese F, Marcotullio MC, Tubaro A, Gianmario A, *et al.* Synthesis and anti-inflammatory activity of natural and semisynthetic geranyloxycoumarins. *Bioorg Med Chem Lett* 2004; 14:2241-2243.
32. Trost BM, Toste FD, Greeman K. Atom economy. Palladium-catalyzed formation of coumarins by addition of phenols and alkynoates via a net C-H insertion. *J Am Chem Soc* 2003; 125:4518-4526.
33. Iranshahi M, Amin GR, Jalalizadeh H, Shafiee A. New germacrane derivative from *Ferula persica* Willd. var *latisecta* Chamberlain. *Pharm Biol* 2003; 41:431-433.
34. Perel'son ME, Sheinker YN, Syrova GP, Turchin KF. NMR spectra of natural coumarin derivatives I. Coumarins. *Chem Nat Comp* 1970; 6:6-14.