

# Effects of pentoxifylline on mouse oocytes maturation and quality *in vitro*

Junjiao Wu<sup>1</sup>, Jianbo Wu<sup>1\*</sup>, Yanyan Xu<sup>1</sup>

<sup>1</sup> Department of obstetrics-gynecology, The First Affiliated Hospital of Fujian Medical University, Fuzhou, 350005, China

## ARTICLE INFO

### Article type:

Original

### Article history:

Received: Feb 3, 2024

Accepted: Jul 31, 2024

### Keywords:

Chromosomes  
Cortical granules  
Oocyte maturation  
Pentoxifylline  
Reactive oxygen species  
Spindle apparatus

## ABSTRACT

**Objective(s):** To investigate the impact of Pentoxifylline (PTX) on the *in vitro* maturation (IVM) of mouse oocytes and its effect on oocyte quality.

**Materials and Methods:** This experimental study involved culturing mouse oocytes in an IVM medium with varying PTX concentrations (0-100  $\mu$ M). Post-culture, oocytes were assessed for nuclear and cytoplasmic maturation and quality indicators, including germinal vesicle breakdown (GVBD), first polar body extrusion (PB1E), cortical granules (CGs) distribution, spindle structure, chromosome alignment, and intracellular reactive oxygen species (ROS) levels.

**Results:** Treatment with PTX at 10, 25, and 50  $\mu$ M concentrations significantly enhanced the nuclear maturation rates of oocytes. The optimal concentration was found to be 10  $\mu$ M, as it resulted in the most favorable cytoplasmic maturation, characterized by improved distribution of CGs, spindle structure, and chromosome alignment. Additionally, treatment with 10  $\mu$ M PTX effectively reduced reactive oxygen species (ROS) levels.

**Conclusion:** PTX supplementation at specific concentrations enhances mouse oocyte maturation and quality, potentially by facilitating CG distribution, spindle integrity, and chromosome alignment and by reducing ROS production.

► Please cite this article as:

Wu J, Wu J, Xu Y. Effects of pentoxifylline on mouse oocytes maturation and quality *in vitro*. Iran J Basic Med Sci 2025; 28:

## Introduction

*In vitro* maturation (IVM) of immature oocytes is a growing topic in Assisted Reproductive Technology (ART), and it has become increasingly popular in treating human infertility in recent years. Experiments have demonstrated, however, that mature oocytes *in vitro* have a much lower developing ability than mature oocytes *in vivo* (1, 3). *In vivo*, the inhibition of nuclear maturation by follicular fluid results in an optimal balance of nuclear and cytoplasmic maturation (4, 5). *In vitro*, the oocyte was removed from the follicular inhibition environment and placed in a culture medium to resume meiosis prematurely without sufficient cytoplasmic maturation (6, 7). As a result of the disorganized nuclear and cytoplasmic maturation, developmental competence is reduced, resulting in a low rate of blastocyst development (8). Therefore, numerous studies expect to increase the cytoplasmic maturation of oocytes by delaying meiosis recovery during IVM (9-11).

Pentoxifylline (PTX) is an old medicine extensively used for treating vascular diseases (12). PTX belongs to the family of non-specific phosphodiesterase (PDE) inhibitors that lead to an increase in cyclic adenosine monophosphate (cAMP) levels, which are known to enhance sperm motility (13, 14). Moreover, it is an anti-oxidant that reduces oxidative stress-induced sperm damage (15, 16).

cAMP has been shown to play a pivotal role in the progression of meiosis in various studies (17-19). PDE

inhibitors elevate cAMP levels, leading to maintenance of meiotic arrest at the diplotene stage and oocyte maturation, ovulation, fertilization, and pregnancy rates (11, 19-21). The reversible inhibitory effects of PDE inhibitors on oocyte maturation and ovulation without altering the menstrual cycle make them promising candidates for non-hormonal therapeutic contraceptive drugs with minimal adverse effects (11). PTX is a member of the PDE inhibitors, although few studies have explored its impact on the *in vitro* maturation of mouse oocytes.

Anti-oxidants regulate ROS, which are necessary for cellular function. Follicles, oocytes, endometrium, and their surroundings are all affected by ROS (22). Overproduction of ROS causes oxidative stress and apoptosis, reducing oocyte quality and developmental potential (23). Therefore, ROS levels are closely linked to oocyte quality. A balance between anti-oxidants and ROS regulates oocyte maturation and quality (24). PTX is an anti-oxidant that reduces the maturation of rat-denuded oocytes cultivated *in vitro*, perhaps by lowering the quantity of ROS (25).

In light of the above, the objective of this present study was to fill a research problem about the impact of PTX on the *in vitro* maturation process of mouse oocytes. Here, we explored the effects of PTX supplementation on the rates of GVBD and PB1E of mouse oocytes during IVM. Additionally, we examined the influence of PTX on the distribution of CGs, spindle structure, chromosome alignment, and intracellular levels of ROS. These findings

\*Corresponding author: Jianbo Wu. Department of obstetrics-gynecology, The First Affiliated Hospital of Fujian Medical University, Fuzhou, 350005, China. Tel: +86-13799395623, Email: gwujianbo@163.com



© 2025. This work is openly licensed via [CC BY 4.0](https://creativecommons.org/licenses/by/3.0/).

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

may elucidate potential mechanisms underlying the beneficial effects of PTX on the maturation and quality of mouse oocytes.

## Materials and Methods

### Mouse and reagents

All animal experiments were performed in strict accordance with the SPF Animal feeding measures of Fujian Medical University's Experimental Animal Research Center and the Ethics Committee (Permit Number: SCXK 2017-0012). Unless otherwise noted, all reagents were purchased from Sigma-Aldrich (USA). ICR mice were housed in a temperature-controlled environment on a 12-hr light/12-hr dark cycle, fed a standard diet, and fed adaptively for one week.

### Oocyte collection and IVM

Female ICR mice aged 5-6 weeks were injected intraperitoneally with 10 IU PMSG and sacrificed 46-48 hr later by cervical dislocation for oocyte collection. Oocytes were collected from ovarian follicles in an immature stage (germinal vesicle) in M2 medium, then treated with hyaluronidase and denuded of cumulus cells using a fine bore glass pipette. Only denuded oocytes with full shape and complete germinal vesicles were used for the following experiments.

M16 medium was used as a maturation medium. Ten to twenty oocytes were cultured in 50  $\mu$ L drops of maturation medium.

### PTX supplementation

PTX was dissolved in the medium at the final concentration of 2.5  $\mu$ M, 5  $\mu$ M, 10  $\mu$ M, 25  $\mu$ M, 50  $\mu$ M, and 100  $\mu$ M and sterilized by filtration.

### Assessment of oocyte maturation by first polar body extrusion

Oocyte maturity was assessed 24 hr after IVM culture, and the oocytes with a first polar body were considered mature under a stereoscopic microscope. The stages of nuclear maturation were GV (germinal vesicle stage), GVBD (when the GV was not visible), and MI (first polar body observed in the perivitelline space). After three replications, the data was analyzed.

### Cortical granule staining

All steps are carried out at room temperature unless otherwise specified.

MII oocytes were washed in DPBS+0.3% PVP and fixed in 4 percent paraformaldehyde for 20 min. The fixed oocytes were removed, exposed to tabletop acid for 20 sec, and washed with CGs blocking solution (DPBS+0.3%BSA+100mM Glycine). The washed oocytes were permeabilized with 1% TritonX-100 dissolved in DPBS for 30 min and then stained for 30 min with FITC-Lens culinary agglutinin (FITC-LCA). Subsequently, oocytes were washed with a CG-blocking solution. Then, DNA was stained by 4, 6-diamidino-2-phenylindole (DAPI) for 10 min. After that, the stained oocytes were mounted on microscope slides visualized under a fluorescence microscope and analyzed by Image-Pro Plus.

### Spindle structure and chromosome staining

MII oocytes were washed in DPBS+0.3% PVP and fixed

with 4% paraformaldehyde for 20 min. The washed oocytes were permeabilized in DPBS for 30 min with 1 percent TritonX-100 dissolved. Oocyte spindles were stained by sequentially incubating the oocytes on monoclonal mouse anti-alpha-tubulin and FITC-goat anti-mouse antibodies. Chromosomes were stained with DAPI and mounted on microscope slides after being thoroughly washed; Image-Pro Plus was used to examine the results.

### Intracellular ROS level

Following 24 hr of IVM, MII oocytes were collected from the control and treatment groups (10  $\mu$ M), washed in DPBS, and incubated in 10 $\mu$ M of DCFH-DA in DPBS for 15 min in the dark. After incubation, the MII oocytes were washed completely in DPBS, placed in a well of a hanging drop slide in 20  $\mu$ L of DPBS, and observed under a fluorescence microscope. Image-Pro Plus was used to evaluate the images of the stained MII oocytes, which were obtained with the same conditions for all groups.

### Statistical analysis

All of the experiments were repeated at least three times. The data in this study were performed using SPSS and Graphpad software. Oocyte maturation in different concentrations was analyzed using one-way ANOVA followed by LSD-t test. Differences in groups of CG distribution, spindle structure, and chromosome alignment were performed with the chi-square. Comparisons of intracellular ROS levels were performed with a t-test. The images presented were analyzed using Image-Pro Plus software. All data was reported as mean values  $\pm$  standard error (SEM), with *P*-values less than 0.05 considered statistically significant.

## Results

### Meiotic progression of mouse oocytes during IVM


We assessed the impact of varying PTX concentrations (0, 2.5, 5, 10, 25, 50, and 100  $\mu$ M) on the meiotic progression of mouse oocytes during IVM. After a 4-hr culture period, the GVBD rate showed a non-significant decrease in the 25  $\mu$ M PTX group compared to other groups (*P*=0.084; Figure 1A, B). Following a 24-hr culture, the PB1E rate was significantly higher in the 10, 25, and 50  $\mu$ M PTX groups than in the control group (*P*<0.01; *P*<0.05; *P*<0.05; Figure 1A, C). In contrast, the 100  $\mu$ M group exhibited fewer oocytes reaching the MII stage.

### Cortical granules distribution

After 24 hr, CG distribution was evaluated under a fluorescence microscope. The groups treated with 2.5, 5, and 10  $\mu$ M PTX displayed higher rates of normal CG distribution, with most CGs migrating to the cortex or forming a cortical granules-free domain (CGFD) (Figure 2B). In contrast, CGs in other groups remained distributed throughout the oocyte cytoplasm (Figure 2A).

### Spindle structure and chromosome alignment

At the conclusion of IVM, spindle structure and chromosome alignment were analyzed (Figure 3A). A reduction in oocytes with dispersed spindles and misaligned chromosomes was observed in the 2.5, 5, 10, and 25  $\mu$ M PTX groups. Notably, the 10  $\mu$ M PTX group showed well-aligned chromosomes and fewer aberrant spindles compared to other groups, suggesting an optimal concentration for



**Figure 1.** Effect of PTX on mouse oocytes IVM (A) After culture for 2, 4, and 24 hr *in vitro*, pictures demonstrated distinct morphologies of mouse oocytes without PTX (control) or with different concentrations of PTX (treatment). Representative oocytes at various stages of development are shown here. The black arrowhead indicates GV, the red arrowhead indicates GVBD, the black arrow indicates MII, and the red arrow indicates matured oocytes with an abnormal first polar body. Scale bar = 100 μm. (B, C) After culture for 4 and 24 hr *in vitro*, bar graphs revealed the percentage of GVBD and PBIE of mouse oocytes. The data is expressed as means ± SEM. \*  $P < 0.05$ , \*\*  $P < 0.01$  versus the control. PTX: Pentoxifylline; IVM: *In vitro* maturation; GV: Germinal vesicle; GVBD: Germinal vesicle breakdown


enhancing cytoplasmic maturation ( $P=0.154$ ,  $P < 0.05$ ; Figure 3B).

**Intracellular ROS level**


The effect of PTX on oxidative stress was determined by measuring intracellular ROS levels post-IVM under a fluorescent microscope. A decrease in ROS signals was noted in oocytes treated with 10 μM PTX, although this did not reach statistical significance ( $P=0.231$ ; Figure 4 A, B).

**Discussion**


PTX is commonly utilized in male ART to assist in the selection of viable sperm and enhance sperm motility prior to intracytoplasmic sperm injection (ICSI) in patients with asthenozoospermia (14, 16, 26). However, little was known about the effect of PTX on mouse oocytes *in vitro*



**Figure 2.** Effect of PTX on CGs distribution of mouse oocytes IVM (A) Representative CGs stained with Lens culinaris agglutinin (LCA) (green) and chromosomes stained with 4, 6-diamidino-2-phenylindole (DAPI) (blue) were shown in the images. The white arrowhead indicates CGFD. Scale bar = 20 μm. (B) Percentage of oocytes having a normal CGs distribution was shown in bar graphs. PTX: Pentoxifylline; CGs: Cortical granules; IVM: *In vitro* maturation



**Figure 3.** Effect of PTX on mouse oocytes spindle structure and chromosome alignment IVM (A) Images showed a representative spindle stained with anti-α-tubulin antibody (green) and chromosomes stained with DAPI (blue). Scale bar = 20 μm. (B) Percentage of oocytes with an abnormal spindle and chromosome was shown in bar graphs. \*  $P < 0.05$  versus the control. PTX: Pentoxifylline; IVM: *In vitro* maturation; DAPI: 4, 6-Diamidino-2-phenylindole



**Figure 4.** Effect of PTX on mouse oocytes intracellular ROS IVM (A) The pictures presented representative oocytes of ROS stained with DCFH-DA (green). Scale bar = 100  $\mu$ m. (B) Fluorescence intensity of ROS per each oocyte was represented by bar graphs. PTX: Pentoxifylline; ROS: Reactive oxygen species; IVM: *In vitro* maturation

maturation. In this study, our primary objective was to investigate whether PTX affects the nuclear maturation of mouse oocytes. Additionally, we explored the influence of PTX on the cytoplasmic maturation of mouse oocytes as part of our secondary investigation. Finally, a series of experiments were conducted to elucidate the underlying mechanism by which PTX promotes *in vitro* maturation of mouse oocytes.

Oocyte maturation is divided into two stages: nuclear and cytoplasmic maturation (27). The stages of nuclear maturation were GV (germinal vesicle stage), GVBD (when the GV was not visible), and MII (first polar body observed in the perivitelline space). We examined the GVBD and PB1E rates in mouse oocytes to focus on whether PTX influenced nuclear maturation (28). According to our findings, the GVBD rate fell only in the 25  $\mu$ M group, while the PB1E rate reduced only in the 100  $\mu$ M group. Additionally, as the concentration of PTX increased in low-concentration groups (2.5, 5, 10), more oocytes of the first polar body were extruded. This contradicts previous studies that specific PDE inhibitors reversibly hinder spontaneous departure from diplotene arrest and oocyte development by raising cyclic nucleotide level (11). According to Chaube's findings, PTX greatly reduced the maturation of denuded rat oocytes, and 1 mM of PTX significantly decreased the rate of maturation ( $P < 0.001$ ) (29). Gupta and Chaube, on the other hand, found that rolipram (PDE 4D inhibitors) did not prevent spontaneous meiotic resumption in denuded oocytes of rats cultivated *in vitro* (10), which was analogous to our findings. These perplexing results might be explained by the fact that PTX has a similar affinity for PDE4 (11, 30) even though PDE4s are plentiful in somatic granulosa cells but not in the oocyte (20, 31). As a result, PDE inhibitors can either inhibit or accelerate oocyte maturation, depending on experimental settings and the species. Consequently, our findings imply that PTX improves nuclear maturation in mouse oocytes at suitable concentrations.

Cytoplasmic maturation is also necessary to allow normal development of embryos (19, 32). The distribution of CGs and the cytoskeleton dynamics are crucial in cytoplasmic maturation and oocyte quality (33, 34). CGs are oocyte cytoplasmic organelles that secrete substances

that prevent polyspermy during acrosome and cortical reactions (35, 36). CGs are limited to the innermost part of the cytoplasm during the GV stage, then migrate to the cortex or form a cortical granules-free domain (CGFD) throughout maturation (37, 38). Moreover, CG migration depends on cytoskeleton function, and orderly meiosis in oocytes necessitates precise control of spindle structure and chromosomal alignment (32, 39). Any mistake in this process can lead to aneuploid eggs, which are the main cause of pregnancy loss and developmental disabilities in humans (40). As a result, mature oocytes should have spindles and chromosomes that are morphologically normal. Here, we discovered that supplementing with PTX during IVM reduced the incidence of aberrant spindle and chromosomal formation while also increasing the rate of normal CG distribution. The findings suggest that PTX supplementation enhances cytoplasmic maturation and quality during *in vitro* oocyte maturation, partly by facilitating normal distribution of CGs, maintaining spindle structure, and promoting chromosome alignment. Based on the findings, the best PTX concentration for mouse oocytes IVM is 10  $\mu$ M. Consequently, the treatment concentration for the following research was 10  $\mu$ M PTX.

Oocyte maturation and quality are regulated by a balance between antioxidants and reactive oxygen species (ROS) (24). The *in vitro* cell culture environment produces a considerable amount of ROS due to the high oxygen concentration than in the *in vivo* environment (41). The physiological concentration of ROS has been demonstrated to play a vital role in normal cell activity, while high concentrations of ROS can be detrimental to cells (42). Excessive ROS production can lead to oxidative stress, disrupting the meiosis process (43). In our study, the rate of aberrant spindles with misaligned chromosomes and abnormal CG distribution was more significant in the control oocytes with elevated ROS levels than in the PTX oocytes supplemented with 10  $\mu$ M. These findings support Premkumar and Chaube's previous research, indicating that PTX effectively mitigates excessive ROS production during *in vitro* oocyte maturation (25).

Throughout our research, at low concentrations (2.5 to 10  $\mu$ M), PTX showed a dose-dependent positive effect, promoting the maturation of oocytes. In contrast, at higher concentrations (25 to 100  $\mu$ M), the effect of PTX showed a dose-dependent negative effect, inhibiting oocyte maturation. Within the concentration range of 2.5 to 10  $\mu$ M, as the concentration of PTX increases, the rates of GVBD and PB1E in oocytes significantly improve, indicating that PTX promotes nuclear maturation of oocytes within this concentration range. However, within the concentration range of 25 to 100  $\mu$ M, the effect of PTX is the opposite. In the 100  $\mu$ M PTX group, there was a decrease in oocytes progressing to the MII stage; concurrently, an increase in the concentration of PTX was correlated with a greater number of abnormal first polar bodies (Figure 1A). This biphasic response may be related to the inhibitory effect of PTX on PDE. PTX promotes oocyte maturation at low concentrations by inhibiting PDE and increasing cAMP levels. At high concentrations, PTX may over-suppress PDE, leading to excessively high levels of cAMP, which interferes with the normal maturation process of oocytes (44, 45). Additionally, high concentrations of PTX may be toxic to cells, affecting cellular structure and function, thereby hindering oocyte maturation. Further research is needed to understand the

mechanisms underlying the observed biphasic response fully and determine the optimal use of PTX in the context of oocyte maturation and fertility treatments.

### Conclusion

In summary, supplementing mouse oocytes with 10  $\mu\text{M}$  PTX significantly improved their maturation and quality *in vitro*. PTX's beneficial effects may be mediated by enhancing CG distribution, spindle structure, and chromosome alignment, which reduces excessive ROS production.

### Acknowledgment

The animals were housed in the Experimental Animal Research Center of Fujian Medical University, China, and other experiments were carried out in the Central Laboratory of Fujian Medical University's First Affiliated Hospital. We express our gratitude to the personnel who work at these centers. The results presented in this paper were part of a student thesis.

### Authors' Contributions

J W supervised, directed, and managed the study; J W designed and conducted the experiments, collected data, and drafted the manuscript; Y X revised the manuscript draft. All authors have reviewed and approved the final version of the manuscript.

### Conflicts of Interest

The authors declare that they have no conflicts of interest.

### References

- LR A. *In vitro* production of embryos in swine. *Theriogenology* 2002; 57:256-273.
- Child TJ, Phillips SJ, Abdul-Jalil AK, Gulekli B, Tan SL. A comparison of *in vitro* maturation and *in vitro* fertilization for women with polycystic ovaries. *Obstet Gynecol* 2002; 100:665-670.
- Eppig JJ, O'Brien MJ, Wigglesworth K, Nicholson A, Zhang W, King BA. Effect of *in vitro* maturation of mouse oocytes on the health and lifespan of adult offspring. *Hum Reprod* 2008; 24:922-928.
- Gosden R, Krapez J, Briggs D. Growth and development of the mammalian oocyte. *Bioassays* 1997; 19:875-882.
- Conti M, Franciosi F. Acquisition of oocyte competence to develop as an embryo: integrated nuclear and cytoplasmic events. *Hum Reprod Update* 2018; 24:245-256.
- Eppig JJ, Schultz RM, O'Brien MJ, Chesnel F. Relationship between the developmental programs controlling nuclear and cytoplasmic maturation of mouse oocytes. *Dev Biol* 1994; 164:1-9.
- Salamone DF, Damiani P, Fissore RA, Robl JM, DUBY RT. Biochemical and developmental evidence that ooplasmic maturation of prepubertal bovine oocytes is compromised. *Biol Reprod* 2001; 64:1761-1768.
- Roy PK, Qamar AY, Tanga BM, Fang X, Kim G, Bang S, *et al.* Enhancing oocyte competence with milrinone as a phosphodiesterase 3A inhibitor to improve the development of porcine cloned embryos. *Front Cell Dev Biol* 2021; 30:1-9.
- Nogueira D, Cortvrindt R, De Matos DG, Vanhoute L, Smits J. Effect of phosphodiesterase type 3 inhibitor on developmental competence of immature mouse oocytes *in vitro*. *Biol Reprod* 2003; 69:2045-2052.
- Gupta A, Chaube SK. Cilostamide and rolipram prevent spontaneous meiotic resumption from diplotene arrest in rat oocytes cultured *in vitro*. *Eur J Pharmacol* 2020; 878:173115.
- Gupta A, Pandey AN, Sharma A, Tiwari M, Yadav PK, Yadav AK, *et al.* Cyclic nucleotide phosphodiesterase inhibitors: possible therapeutic drugs for female fertility regulation. *Eur J Pharmacol* 2020; 883:1-6.
- Donate-Correa J, Tagua VG, Ferri C, Martín-Núñez E, Hernández-Carballo C, Ureña-Torres P, *et al.* Pentoxifylline for renal protection in diabetic kidney disease. A model of old drugs for new horizons. *J Clin Med* 2019; 8:287-303.
- Shen MR, Chiang PH, Yang RC, Hong CY, Chen SS. Pentoxifylline stimulates human sperm motility both *in vitro* and after oral therapy. *Brit J Clin Pharmacol* 1991; 31:711-714.
- Xian Y, Jiang M, Liu B, Zhao W, Zhou B, Liu X, *et al.* A cryoprotectant supplemented with pentoxifylline can improve the effect of freezing on the motility of human testicular sperm. *Zygote* 2021; 22:1-6.
- Dhulqarnain AO, Takzaree N, Hassanzadeh G, Tooli H, Malekzadeh M, Khanmohammadi N, *et al.* Pentoxifylline improves the survival of spermatogenic cells via oxidative stress suppression and upregulation of PI3K/AKT pathway in mouse model of testicular torsion-detorsion. *Helion* 2021; 7:1-11.
- Beygi Z, Forouhari S, Mahmoudi E, Hayat SMG, Nourimand F. Role of oxidative stress and antioxidant supplementation in male fertility. *Curr Mol Med* 2021; 21:265-282.
- S B. Cows are not mice: the role of cyclic AMP, phosphodiesterases, and adenosine monophosphate-activated protein kinase in the maintenance of meiotic arrest in bovine oocytes. *Mol Reprod Dev* 2011; 78:734-742.
- Mermillod P, Tomanek M, Marchal R, Meijer L. High developmental competence of cattle oocytes maintained at the germinal vesicle stage for 14 hr in culture by specific inhibition of MPF kinase activity. *Mol Reprod Dev* 2000; 55:89-95.
- Ramos Leal C, Santos Monteiro CA, Souza-Fabjan JMG, de Paula Vasconcelos CO, Garcia Nogueira LA, Reis Ferreira AM, *et al.* Role of cAMP modulator supplementations during oocyte *in vitro* maturation in domestic animals. *Anim Reprod Sci* 2018; 199:1-14.
- Conti M. Phosphodiesterases and regulation of female reproductive function. *Curr Opin Pharmacol* 2011; 11:665-669.
- Gur a A, Tiwari M, Prasad S, Chaube SK. Role of cyclic nucleotide phosphodiesterases during meiotic resumption from diplotene arrest in mammalian oocytes. *J Cell Biochem* 2017; 118:446-452.
- Kala M, Shaikh MV, Nivsarkar M. Equilibrium between anti-oxidants and reactive oxygen species: a requisite for oocyte development and maturation. *Reprod Med Biol* 2017; 16:28-35.
- Dai X, Lu Y, Zhang M, Miao Y, Zhou C, Cui Z, *et al.* Melatonin improves the fertilization ability of post-ovulatory aged mouse oocytes by stabilizing ovastacin and juno to promote sperm binding and fusion. *Hum Reprod* 2017; 2017:1-30.
- Wang S, He G, Chen M, Zuo T, Xu W, Liu X. The role of antioxidant enzymes in the ovaries. *Oxid Med Cell Longev* 2017; 2017:1-14.
- Premkumar KV, Chaube SK. Increased level of reactive oxygen species persuades postovulatory aging-mediated spontaneous egg activation in rat eggs cultured *in vitro*. *In Vitro Cell Dev Biol Anim* 2016; 52:576-588.
- Satish M, Kumari S, Deeksha W, Abhishek S, Nitin K, Adiga SK, *et al.* Structure-based redesigning of pentoxifylline analogs against selective phosphodiesterases to modulate sperm functional competence for assisted reproductive technologies. *Sci Rep* 2021; 11:12293-12310.
- Ferrer-Vaquero A, Barragán M, Rodríguez A, Vassena R. Altered cytoplasmic maturation in rescued *in vitro* matured oocytes. *Hum Reprod* 2019; 34:1095-1105.
- Hoshino Y. Updating the markers for oocyte quality evaluation: intracellular temperature as a new index. *Reprod Med Biol* 2018; 17:434-441.
- Chaube SK. Effects of pentoxifylline and caffeine on spontaneous maturation of rat oocytes. *Health and Population; Perspectives and Issues* 2000; 23:177-189.

30. Yougbare I, Belemnaba L, Morin C, Abusnina A, Senouvo YF, Keravis T, *et al.* NCS 613, a potent PDE4 inhibitor, displays anti-inflammatory and anti-proliferative properties on A549 lung epithelial cells and human lung adenocarcinoma explants. *Front Pharmacol* 2020; 11:1266-1275.
31. Tsafiriri A, Chun S, Zhang R, Hsueh AJW, Conti M. Oocyte maturation involves compartmentalization and opposing changes of cAMP levels in follicular somatic and germ cells: studies using selective phosphodiesterase inhibitors. *Dev Biol* 1996; 178:393-402.
32. Park JE, Kim M, Lee J, Chai C, Lee ST, Lee E. *in vitro* maturation on an agarose matrix improves the developmental competence of porcine oocytes. *Theriogenology* 2020; 157:7-17.
33. Lin T, Lee JE, Kang JW, Oqani RK, Cho ES, Kim SB, *et al.* Melatonin supplementation during prolonged *in vitro* maturation improves the quality and development of poor-quality porcine oocytes via anti-oxidative and anti-apoptotic effects. *Mol Reprod Dev* 2018; 85:665-681.
34. Lin T, Lee JE, Oqani RK, Kim SY, Cho ES, Jeong YD, *et al.* Tauroursodeoxycholic acid improves pre-implantation development of porcine SCNT embryo by endoplasmic reticulum stress inhibition. *Reprod Biol* 2016; 16:269-278.
35. Jeseta M, Budna J, Kranc W, Hanulakova S, Bryja A, Chachuła A, *et al.* Expression of genes encoding zona pellucida glycoproteins and cortical granule distribution in porcine oocytes isolated from small and medium follicles in relation to puberty status of donors. *Czech J Anim Sci* 2017; 62:234-241.
36. Vogt E, Tokuhiko K, Guo M, Dale R, Yang G, Shin S, *et al.* Anchoring cortical granules in the cortex ensures trafficking to the plasma membrane for post-fertilization exocytosis. *Nat Commun* 2019; 10:2271-2284.
37. Coticchio G, Dal Canto M, Fadini R, Mignini Renzini M, Guglielmo MC, Miglietta S, *et al.* Ultrastructure of human oocytes after *in vitro* maturation. *Mol Hum Reprod* 2016; 22:110-118.
38. Liu X, Mal S, Miao D, Liu D, Bao S, Tan J. Cortical granules behave differently in mouse oocytes matured under different conditions. *Hum Reprod* 2005; 20:3402-3413.
39. Ma R, Zhang J, Liu X, Li L, Liu H, Rui R, *et al.* Involvement of Rab6a in organelle rearrangement and cytoskeletal organization during mouse oocyte maturation. *Sci Rep* 2016; 6:23560-23569.
40. Hassold T, Hunt P. To err (meiotically) is human: the genesis of human aneuploidy. *Nat Rev Genet* 2001; 2:280-291.
41. Cetica PD, Pintos LN, Dalvit GC, Beconi MT. Antioxidant enzyme activity and oxidative stress in bovine oocyte *in vitro* maturation. *IUBMB Life* 2001; 51:57-61.
42. Hancock JT, Desikan R, Neill SJ. Role of reactive oxygen species in cell signalling pathways. *Biochem Soc Trans* 2001; 29:345-350.
43. Di Emidio G, Falone S, Vitti M, D'Alessandro AM, Vento M, Di Pietro C, *et al.* SIRT1 signalling protects mouse oocytes against oxidative stress and is deregulated during aging. *Hum Reprod* 2014; 29:2006-2017.
44. Jaffe LA, Egbert JP. Regulation of mammalian oocyte meiosis by intercellular communication within the ovarian follicle. *Annu Rev Physiol* 2007; 69:237-260.
45. Zhang M, Su Y, Sugiura K, Xia G, Eppig JJ. Granulosa cell ligand LPPC and its receptor NPR2 maintain meiotic arrest in mouse oocytes. *Science* 2010; 330:366-369.