

Role of gallic acid in sarcopenia in D-galactose-treated mice

Xianchu Liu ^{1*}, Zile Zhu ¹, Ming Liu ², Zhihe Mao ^{1*}

¹ Institute for Scientific Sports and Health Promotion, Hunan University of Arts and Science, Changde, Hunan, China

² Faculty of Science, College of Furong, Hunan University of Arts and Science, Changde, Hunan, China

ARTICLE INFO

Article type:

Original

Article history:

Received: Aug 1, 2025

Accepted: May 23, 2026

Keywords:

Aging

Apoptosis

Autophagy

Gallic acid

Mitochondrion

Sarcopenia

ABSTRACT

Objective(s): Aging is a natural phenomenon that results in sarcopenia to impair living quality in humans. Gallic Acid (GA) has many biological activities against aging and muscle injury. Our purpose was to clarify possible mechanisms of GA on aged-relevant sarcopenia.

Materials and Methods: Senescent model was established by D-galactose in mice. The senescent model was treated with GA for 10 weeks. Skeletal muscle was stained via HE to assess muscle fiber size. Oxidation indicators and inflammatory factors were detected using a spectrophotometer and ELISA. The protein expressions were tested by Western blot.

Results: Our data unveiled GA enhanced muscle mass to improve grip strength and fatigue in D-galactose-induced aging mice. In serum, GA increased T-AOC, SOD and CAT levels, and decreased MDA, TNF- α , and IL-6 levels to exert its anti-oxidative and anti-inflammatory properties. In skeletal muscle, GA decreased atrogen 1, Murf-1, collagen I, and Tgf- β 1 expressions to improve atrophy and fibrosis. Furthermore, GA increased Pgc-1 α , Nrf-1, Bcl-2, and Bcl-2 expressions, and decreased P62 and Bax expressions to prevent skeletal muscle damage.

Conclusion: GA exerted its protective effect on sarcopenia concomitant aging via moderating mitochondrial biology, enhancing autophagy and restraining apoptosis.

► Please cite this article as:

Liu X, Zhu Z, Liu M, Mao Z. Role of gallic acid in sarcopenia in D-galactose-treated mice. Iran J Basic Med Sci 2026; 29:

Introduction

Sarcopenia is one of age-related degenerative conditions, which is also related to multiple co-morbidities, such as cognitive impairment and diabetes mellitus (1, 2). Skeletal muscle is a primary tissue responsible for movement, whole-body metabolism and overall health. With increasing age, skeletal muscle gradually degenerates and motor capacity displays a downwards trend (3). Sarcopenia is linked to a reduction in muscle weight and grip strength, which influence life quality and athletic ability (4). These alterations are closely connected with the functional state of skeletal muscles. Sarcopenia leads to increased risk of falls and disability. In addition, the problem of aging is getting worse year by year. The proportion of elderly people will reach one in six individuals in 2050. Accordingly, the morbidity of aged-relevant sarcopenia increases with time (5). However, the feasible therapeutic method underlying aged-relevant sarcopenia remains unclear.

There are various kinds of factors that may result in sarcopenia. One of the reasons for aged-relevant sarcopenia is skeletal muscle dystrophy caused by ubiquitin-proteasome system disorders (6). Ubiquitin-mediated proteolysis plays a crucial role in muscle atrophy (7). Disturbance of proteostasis intensifies proteolysis to lessen muscle weight (8). Atrogen 1 and Murf-1 are crucial ubiquitin ligases, which are involved in regulating protein homeostasis.

Ubiquitin-proteasome system is activated as represented by increased atrogen 1 and Murf-1 to induce muscle proteolysis (9). Fibrosis is another vital pathological characteristic of skeletal muscle aging, which accelerates aged-relevant sarcopenia process (10). Collagen I and Tgf- β 1 as important regulatory factors for fibrosis are activated in various organs of the elderly (11). Likewise, advancing age is proved to be connected with a condition of fibrosis in skeletal muscles, which is characterized by regulating the TGF- β pathway. Hence, suppression of muscle atrophy and fibrosis may be an effective way in the treatment of aged-relevant sarcopenia.

Gallic Acid (GA) is a plant polyphenolic compound and can be extracted from natural plants, such as strawberries and red wines. Recently, GA has been applied in medicine and functional food because of its extensive biological function, such as anti-oxidative stress potential and anti-inflammatory effects (12, 13). Several researches revealed GA possessed anti-aging effect in many experiments. The Alzheimer model research revealed GA as a dual α/β -secretase modulator mitigated cerebral amyloidosis and reversed senescence-associated cognitive impairment (14). Another research demonstrated GA regulated senescence-related molecular and biochemical indicators (15). GA was uncovered to be strongly associated with mitigation of d-galactose induced thymic involution via its anti-apoptosis capacity (16). Recent research has suggested GA alleviated

*Corresponding authors: Liu Xianchu. Institute for Scientific Sports and Health Promotion, Hunan University of Arts and Science, Changde, Hunan, China. Email: liuxc645@huas.edu.cn; Mao Zhihe. Institute for Scientific Sports and Health Promotion, Hunan University of Arts and Science, Changde, Hunan, China. Email: maozhihe1729@huas.edu.cn



© 2026. This work is openly licensed via [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/).

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

D-galactose-induced cardiac hypertrophy via modulation of mitochondrial biogenesis (17). However, the anti-aging effect of GA on skeletal muscles is not expounded. Therefore, we hypothesized GA might play a vital in improvement of muscle atrophy and fibrosis during aging, and explored its biologic activity and mechanism in the development of sarcopenia in D-galactose-induced aged mice.

Materials and Methods

Animals

Forty male ICR mice were 6-8 weeks old. The body weight of mice was 18-22 g. The experiments were approved by Ethics Committee of Hunan University of Arts and Science, China (No. HUAS-2021-TY-085).

Experimental design

Mice were divided into 4 groups. Each group included 10 mice. The Control group was administered normal saline solution. Gallic acid (Purity \geq 99.0%) was purchased from Sangon Biotech. GA was stored at 4 °C. The Model group was subcutaneously injected with D-galactose (18). The injected dose of D-galactose was 500 mg/kg body weight. The GA group received D-galactose and gallic acid. Experimental animals of GA-L group and GA-H group were administered intragastric GA at different doses for 10 weeks. The dose of GA was 50 or 100 mg/kg/d.

Grip strength test

Mice pulled stick to test grip strength. Grip tester was purchased from Anhui Zhenghua Bioinstrumentation. Grip strength of experiment data was acquired by multiple metering.

Forced swimming test

After grip strength test, mice underwent a loading swimming. The water depth was 30 cm. For measuring exhaustive time, the mouse was forced to swim with load (19). Exhaustive time of each mouse was calculated when experimental animal could not swim over 10 sec.

Histological analysis

After forced swimming test, animals were sacrificed. Gastrocnemius and soleus were dissected and weighed. Gastrocnemius was fixed, paraffin embedded and cut into 4 μ m (20). The nucleus and cytoplasm of gastrocnemius stained with Hematoxylin and Eosin (HE), respectively. The image was gathered through an optical microscope. The cross-sectional area of gastrocnemius tissue was counted via ImageJ.

Biochemical assay

Blood was acquired from the eyeball. Then, blood was centrifuged to separate the supernatant. The detection kits of biochemical indicators were purchased from Nanjing Jiancheng Biotechnology Institute. The absorbance values of ALT, AST, BUN, and CK were detected at 505, 510, 640, and 660 nm by a spectrophotometer.

Oxidation indices assay

The detection kits of oxidation indices were purchased from Nanjing Jiancheng Biotechnology Institute. Gastrocnemius was ground and centrifuged. The supernatant was used to assay oxidation indices. SOD, CAT, and MDA

were determined by the hydroxylamine method, ammonium molybdate spectrometric method, and thibaburic acid method. The absorbance values of CAT, MDA, SOD, and T-AOC were detected at 405, 532, 550, and 520 nm.

ELISA

The inflammatory cytokines were detected by ELISA. The ELISA kits were purchased from BOSTER Biological Technology. Tetramethylbenzidine was used as chromogenic substrate. The absorbance value was tested at 405 nm.

Western blot

Skeletal muscle was rapidly frozen by liquid nitrogen. Protein was stored at -80 °C. Different molecular weights of protein were separated by SDS-PAGE. The expression of proteins was tested via western blot. The protein of skeletal muscle was transferred onto PVDF by wet electroblot technology. The antibodies of collagen I (1:500 dilution), Tgf- β 1 (1:1000 dilution), BAX (1:1000 dilution), BCL-2(1:1000 dilution), β -actin (1:1000 dilution), and HRP-conjugated anti-mouse (1:3000 dilution) and anti-rabbit (1:3000 dilution) were obtained from Proteintech (Wuhan, China). The antibodies of Troponin-1(1:500 dilution) and MuRF-1(1:500 dilution), NF- κ B(1:500 dilution), and PGC-1 α (1:500 dilution) were obtained from Sangon Biotech (Shanghai, China). The antibodies of Beclin-1 (1:500 dilution) and P62 (1:1000 dilution) were obtained from Servicebio (Wuhan, China). β -actin gene was used as internal reference. Protein signal was captured by ECL system.

Statistical analysis

All results were expressed as Mean \pm SD. The significance was demonstrated by a one-way ANOVA followed by Duncan's Multiple Range test. $P < 0.05$ was deemed significant.

Results

Role of GA in skeletal muscle dysfunction

Sarcopenia has a high incidence in the aged adults. Aged-relevant sarcopenia impairs fragility and mobility in skeletal muscles. The dominant characteristic of sarcopenia is muscle loss which leads to decreased muscle strength. In Figure 1A-B, D-gal obviously decreased myocyte cross-sectional area ($P < 0.01$), while GA visibly augmented myofiber sizes ($P < 0.05$). In Figure 1C-D, D-gal obviously decreased the weight of gastrocnemius and soleus ($P < 0.01$), while GA visibly increased muscle mass ($P < 0.05$). In Figure 1E-F, D-gal obviously reduced grip strength and exercise performance, while GA visibly increased muscle function ($P < 0.01$).

Role of GA in biochemical indexes in serum

Aging is a complex disease with various premonitory symptoms which is relevant to physiological dysfunctions in many organisms. D-galactose is widely used to mimic natural aging in human. In animal experiments, D-galactose was proved to cause multiple organism injuries. In Figure 2, D-gal obviously induced CK, BUN, ALT, and AST levels ($P < 0.01$), while GA visibly decreased above biochemical indicators ($P < 0.01$).

Role of GA in oxidation injury in serum

An increase in oxidative stress is a common characteristic during the aging process. Antioxidant enzymes can facilitate




Figure 1. Role of gallic acid (GA) in skeletal muscle dysfunction in senescent mice (A) Representative images of gastrocnemius stained with H&E (200X magnification), (B) muscle fiber size, (C) gastrocnemius weight, (D) soleus weight, (E) grip strength, and (F) exhaustion time. ** $P < 0.01$ vs control. * $P < 0.05$ vs Model. ## $P < 0.01$ vs model

oxidation resistance to increase cell survival by eliminating reactive oxygen species. Accumulation of lipid peroxidation is demonstrated to aggravate oxidative damage. In Figure 3, D-gal obviously reduced T-AOC, SOD and CAT contents ($P < 0.01$), while MDA concentration was obviously elevated ($P < 0.01$). However, GA possibly alleviated oxidation injury, to sustain internal environment homeostasis ($P < 0.01$).




Figure 2. Role of gallic acid (GA) in biochemical indexes in senescent mice Serum levels of (A) CK, (B) BUN, (C) ALT, and (D) AST. ** $P < 0.01$ vs Control. ## $P < 0.01$ vs Model CK: Creatine kinase; BUN: Blood urea nitrogen; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase




Figure 3. Role of gallic acid (GA) in oxidation injury in senescent mice Serum contents of (A) T-AOC, (B) SOD, (C) CAT, and (D) MDA. ** $P < 0.01$ vs Control. ## $P < 0.01$ vs Model T-AOC: Total antioxidant capacity; SOD: Super oxide dismutase; CAT: Catalase; MDA: Malondialdehyde

Role of GA in inflammatory response in serum

Inflammatory response is demonstrated as a prominent activation during the aging process, which is taken for contributing to organism damage. Aging results in the overproductions of TNF- α and IL-6 in the body. In Figure 4, D-gal obviously promoted TNF- α and IL-6 concentrations ($P < 0.01$). However, GA possibly suppressed inflammatory reaction ($P < 0.01$).

Role of GA in atrophy in skeletal muscle

The aged-induced muscle loss is a result of protein degradation. Skeletal muscle atrophy is a crucial pathological change in sarcopenia. The ubiquitin-proteasome system plays a major role in muscle atrophy. In Figure 5, D-gal obviously strengthened atrogen 1 and Murf-1 expressions ($P < 0.01$). However, GA possibly alleviated D-galactose-evoked increase of protein degeneration factors to improve muscle atrophy ($P < 0.01$).

Role of GA in fibrosis in skeletal muscle

In sarcopenia, aging aggravates myopathy and muscle structure change, as represented by accelerating fibrosis in skeletal muscles. Collagen I is a marker factor to exhibit collagen fibers deposition and fibrotic changes. Tgf- β 1 is a potent profibrogenic cytokine and can activate fibrosis process in skeletal muscles. In Figure 6, D-gal obviously enhanced collagen I and Tgf- β 1 expressions ($P < 0.01$). However, GA possibly restrained the TGF- β pathway to improve muscle fibrosis ($P < 0.01$).

Role of GA in mitochondrial disorder in skeletal muscle

Mitochondrial disorder is an important inducement during the aging process that aggravates the development of sarcopenia. Mitochondrion is a major site of energy production, which plays a vital role in exercise performance. Pgc-1 α as a master regulator of mitochondrion is involved in activating Nrf-1. In Figure 7, D-gal obviously reduced Nrf-1 and Pgc-1 α expressions ($P < 0.01$). However, GA possibly mitigated D-galactose-evoked mitochondrial disorder in skeletal muscle ($P < 0.01$).




Figure 4. Role of gallic acid (GA) in inflammatory response in senescent mice. Serum concentrations of (A) TNF- α and (B) IL-6. ** $P < 0.01$ vs Control. ## $P < 0.01$ vs Model




Figure 5. Role of gallic acid (GA) in muscle atrophy in senescent mice. Western blot and semiquantitative analysis of (A, B) atrogen 1 and (C, D) Murf-1. ** $P < 0.01$ vs Control. ## $P < 0.01$ vs Model




Figure 5. Role of gallic acid (GA) in mitochondrial disorder in senescent mice. Western blot and semiquantitative analysis of (A, B) Nrf-1 and Pgc-1 α (C, D). ** $P < 0.01$ vs Control. ## $P < 0.01$ vs Model




Figure 6. Role of gallic acid (GA) in muscle fibrosis in senescent mice. Western blot and semiquantitative analysis of (A, B) collagen I and (C, D) Tgf- β 1. ** $P < 0.01$ vs Control. ## $P < 0.01$ vs Model




Figure 8. Role of gallic acid (GA) in autophagy in senescent mice. Western blot and semiquantitative analysis of (A, B) Beclin-1 and (C, D) P62. ** $P < 0.01$ vs Control. ## $P < 0.01$ vs Model

Role of GA in autophagy in skeletal muscle

Autophagy plays a key role in aging and muscular disorders which is closely associated with sarcopenia. Autophagy is part of catabolic pathways and can clear away damaged organelles via lysosomal machinery to improve physiology. In Figure 8, D-gal obviously restrained Beclin-1 expression and heightened P62 expression ($P < 0.01$). However, GA possibly relieved D-galactose-evoked abnormal autophagy in skeletal muscles ($P < 0.01$).

Role of GA in apoptosis in skeletal muscles

Apoptosis pathway is visibility activated during the aging process, which is believed to results in sarcopenia. The ratio of Bax/Bcl-2, as integral parts in apoptotic cascade, is credibly involved in playing an important role in moderation of apoptotic machinery. In Figure 9, D-gal obviously heightened Bax expression and restrained Bcl-2 expression ($P < 0.01$). However, GA possibly alleviated D-galactose-evoked excessive apoptosis in skeletal muscle ($P < 0.01$).




Figure 9. Role of gallic acid (GA) in apoptosis in senescent mice. Western blot and semiquantitative analysis of (A, B) Bax and (C, D) Bcl-2. ** $P < 0.01$ vs Control, # $P < 0.01$ vs model

Discussion

Sarcopenia is associated with aging. The symptom of sarcopenia is muscle loss (21). Increased with age, less muscle mass affects skeletal muscle function (22). Previous research showed GA possessed a variety of physiological effects on skeletal muscle. GA could be used as a skeletal muscle relaxant in animal experiment (23). In high-fat-fed mice, GA partially reduced plasma FFA and intramuscular lipid deposition (24). In streptozotocin-treated rats, GA partially stimulated insulin secretion and enhanced GLUT4 expression in soleus muscle tissue to improve diabetes mellitus (25). In type 2 diabetic rats, GA promoted glucose uptake via PI3K/p-Akt signaling pathway and slightly enhanced PPAR γ expressions in skeletal muscles (26). D-galactose is an inducer of sarcopenia. During aging, skeletal muscle weight is reduced and athletic ability is weakened. In this study, GA alleviated D-galactose-induced skeletal muscle loss, muscle force decline and dyskinesia, suggesting GA might be involved in ameliorating sarcopenia in aged mice.

Aging is a harm factor of healthy longevity and causes many organs senescent, such as muscle, kidney, and liver. Previous research showed GA had an extensive protective effect on muscles, liver and kidney injury. In snake envenoming explored rat, GA partially reduced CK activity to lessen muscle damage and increase myogenin to accelerate muscle repair (27). In sodium arsenite-induced renal toxicity, GA decreased BUN levels to relieve kidney damage (28). In carbon tetrachloride-induced liver toxicity, GA decreased ALT and AST levels to relieve hepatic damage (29). D-galactose can induce increased CK, BUN, ALT, and AST levels during aging. In this study, GA reversed above adverse biochemical indicators levels in aged mice, suggesting GA might ameliorate muscle, kidney, and hepatic damage against D-galactose-induced aging.

The mechanism of aging is complex. Oxidative imbalance is a vital incentive of aging, which plays a primary role in oxidative stress-induced geriatric syndromes. During aging, oxidative products are produced in large quantities, while oxidation resistance is reduced in the living organism. Previous research showed GA protected living organisms from peroxidation. GA affected carbohydrate metabolism and enhanced T-AOC level to alleviate oxidative stress (30). In cyclophosphamide-treated mice, GA enhanced CAT

activity to retard renal toxicity (31). In ethanol exposed zebrafish, GA degraded extracellular nucleotides and enhanced SOD activity to suppress brain damage (32). In hyperuricemic mice, GA regulated urate transporters and relieved MDA to suppress kidney damage (33). The old people are susceptible to be impaired by ROS because of the reduced capacity of anti-oxidization. During aging, D-galactose induced decrease of T-AOC and increase of MDA levels in serum. In addition, antioxidant enzymes levels might be decreased in senescent mice. In this study, GA recovered oxidation resistance and reduced oxidative products in aged mice, suggesting GA might ameliorate internal oxidation against D-galactose-induced aging.

Aging aggravates inflammatory response to cause organism damage. Previous research showed GA possessed anti-inflammatory effects in many disease models. GA decreased inflammatory cytokine levels in serum to mitigate pain and depression (34). In non-alcoholic fatty mice, GA repressed inflammatory pathways to protect liver functions (35). In cigarette smoke-treated mice, GA blunted TNF- α expression to protect pulmonary inflammation (36). Up-regulation of TNF- α and IL-6 caused an imbalance in inflammatory response. In addition, D-galactose induced TNF- α and IL-6 levels. In this study, GA reduced TNF- α and IL-6 levels in aged mice, suggesting GA might ameliorate inflammatory response against senility.

Muscle atrophy is the main cause of sarcopenia. UPS is involved in proteolysis to regulate protein homeostasis and muscle mass. In other words, loss of proteostasis leads to muscle atrophy. Up-regulation of ubiquitin ligase caused a disorder in ubiquitin-proteasome system. In STZ diabetes, GA inhibited atrogen 1 expression in skeletal muscle to prevent myopathy (37). In C2C12 cells, GA partially inhibited Murf-1 expression to promote skeletal muscle development (38). In addition, D-galactose increased atrogen 1 and Murf-1 expressions in skeletal muscles to induce sarcopenia. In this study, GA reduced atrogen 1 and Murf-1 expressions in aged mice, suggesting GA might sustain the ubiquitin-proteasome system against D-galactose-induced sarcopenia.

Fibrosis is a crucial pathological alteration in musculoskeletal system disease. D-galactose aggravated skeletal muscle injury, which was involved in fibrosis. In transverse aortic constriction-treated mice, GA suppressed connective tissue growth factor and deposition of collagen I to mitigate cardiac fibroblasts (39). In type I diabetes rats, GA possibly inhibited fibronectin expression to positively affect testicular fibroblasts (40). In glyoxal-treated rats, GA suppressed collagen deposition and collagen III mRNA expression to mitigate renal fibrosis (41). In thioacetamide-treated rats, GA was involved in regulating the Tgf- β 1/Smad3 pathway to exhibit hepatoprotective and mitigate liver fibrosis (42). In bleomycin-treated mice, GA modulated the Tgf- β 1/Smad2 pathway and balanced NOX4/Nrf2 to attenuate pulmonary fibrosis (43). In addition, D-galactose increased collagen I and Tgf- β 1 expressions to induce skeletal muscle fibrosis. In this study, GA might reduce TGF- β pathway against senility.

Mitochondrial dysfunction is nearly related to all geriatric diseases. Mitochondria, as primary sites of energy production, play an important role in athletic ability. During the aging process, mitochondrial dysfunction is more and more prominent. In obese mice, GA enhanced glucose

tolerance and increased PGC1- α mRNA expression in brown adipose tissue (44). In bisphenol A-exposed rats, GA was involved in improving enzymes of respiratory chain and intactness of mitochondria and pyruvate dehydrogenase to sustain number and structure of mitochondria against liver injury (45). In Na₂S₂O₄-exposed SH-SY5Y cells, GA improved mitochondrial membrane potential, oxygen consumption, and ATP synthesis levels in hypoxia/reoxygenation-induced mitochondrial discords (46). In aging rats, GA regulated Pgc-1 α , TFAM and Nrf2 mRNA expressions to protect mitochondria against D-galactose-induced heart injury (17). Pgc-1 α plays an important role in mitochondrial oxidative metabolism. Nrf-1 is involved in modifying mitochondrial function. In skeletal muscles, D-galactose decreased Nrf-1 and Pgc-1 α in aged mice. In this study, GA prevents these changes induced by D-galactose in aged mice, suggesting GA might regulate mitochondrial biology against aged-relevant sarcopenia.

Autophagy pathway in skeletal muscles is associated with sarcopenia. During the aging process, autophagy was inhibited to cause many kinds of organ injury, including skeletal muscles. In addition, autophagy, as a vital proteolytic cascade, is involved in cleaning injured mitochondria to maintain skeletal muscle structure and function. Regulating autophagy might be a viable therapeutic approach to improve geriatric diseases. GA enhanced autophagic flux, as indicated by decreasing p62 protein levels in liver (47). In TAC-induced mice, GA activated the autophagy-dependent mechanism via the activation of Beclin-1 protein levels to improve cardiac hypertrophy and myocardial fibrosis (48). In ischemia-reperfusion rats, GA regulated Beclin-1 expression to protect the intestinal tissue (49). In skeletal muscles, D-galactose increased Beclin-1 and decreased P62 expressions in aged mice. In this study, GA prevents these changes induced by D-galactose in aged mice, suggesting GA might regulate autophagy pathway against aged-relevant sarcopenia.

Apoptosis dysbiosis is closely related to muscle damage, which leads to muscle force decline and dysfunction. During the aging process, aging induced apoptosis dysbiosis to affect normal physiological functions in the body. In skeletal muscles, aging aggravates apoptosis to cause muscle loss. In addition, apoptosis is linked to mitochondrial function and autophagic flux. In spontaneously hypertensive rats, GA prevented S mRNA levels to relieve CaMKII δ -induced apoptosis (50). In hypothyroid rats, GA suppressed oxidative, endoplasmic reticulum stresses and Bax/Bcl-2 ratio to attenuate adult-onset hippocampal damage (51). In type 2 diabetic rats, GA possibly ameliorated glucose tolerance and enhanced Bcl-2 expression in hippocampus to improve neurodegeneration (52). In aging rats, GA modulated apoptosis against D-galactose-induced cardiac hypertrophy (17). In skeletal muscles, D-galactose mitigated Bax and accelerated Bcl-2 expressions in aged mice. In this study, GA prevents these changes induced by D-galactose in aged mice, suggesting GA might regulate apoptosis pathway against aged-relevant sarcopenia.

Conclusion

In this study, GA was proved to ameliorate muscle mass and weight to enhance grip strength and athletic ability in aged mice induced by D-galactose. In serum, GA lowered oxidative stress, enhanced antioxidant status, and

diminished inflammation, which contribute to its possible anti-aging function induced by D-galactose. In skeletal muscles, GA improved atrophy and fibrosis, making it possibly beneficial to preventing D-galactose-induced sarcopenia. Mechanistically, the roles of GA were related to its functions in amelioration of mitochondrial biology, enhancement of autophagy, and restraint of apoptosis. Therefore, GA could be used to improve sarcopenia because of its theoretical basis in anti-aging and muscular protection.

Acknowledgment

This study was supported in part by Hunan Provincial Natural Science Foundation of China (Grant No. 2024JJ5294), Scientific Research Fund of Hunan Provincial Education Department (Grant No. 23A0506), and Changde City's Guiding Projects for Technological Innovation (Grant No. 2024ZD43).

Authors' Contributions

L X contributed to study design. L X, Z Z, M Z, and L M carried out experiments. L X and M Z discussed the results. L X and L M drafted the manuscript and revised the final manuscript critically.

Conflicts of Interest

The authors declare no conflicts of interest.

Declaration

We have not used any AI tools or technologies to prepare this manuscript.

References

1. Peball M, Mahlkecht P, Werkmann M, Marini K, Murr F, Herzmann H, *et al.* Prevalence and associated factors of sarcopenia and frailty in Parkinson's disease: A cross-sectional study. *Gerontology* 2019; 65: 216-228.
2. Souza ABF, Nascimento DAC, Rodrigues IJM, Charone CCO, Lopes GL, Lima RS, *et al.* Association between sarcopenia and diabetes in community dwelling elderly in the Amazon region - Viver Mais Project. *Arch Gerontol Geriatr* 2019; 83: 121-125.
3. Tieland M, Trouwborst I, Clark BC. Skeletal muscle performance and ageing. *J Cachexia Sarcopenia Muscle* 2018; 9: 3-19.
4. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, *et al.* Sarcopenia: European consensus on definition and diagnosis: Report of the European working group on sarcopenia in older people. *Age Ageing* 2010; 39: 412-423.
5. Wilkinson DJ, Piasecki M, Atherton PJ. The age-related loss of skeletal muscle mass and function: Measurement and physiology of muscle fibre atrophy and muscle fibre loss in humans. *Ageing Res Rev* 2018; 47: 123-132.
6. Yin L, Li N, Jia W, Wang N, Liang M, Yang X, *et al.* Skeletal muscle atrophy: From mechanisms to treatments. *Pharmacol Res* 2021; 172: 105807.
7. Sinam IS, Chanda D, Thoudam T, Kim MJ, Kim BG, Kang HJ, *et al.* Pyruvate dehydrogenase kinase 4 promotes ubiquitin-proteasome system-dependent muscle atrophy. *J Cachexia Sarcopenia Muscle* 2022; 13: 3122-3136.
8. Bilodeau PA, Coyne ES, Wing SS. The ubiquitin proteasome system in atrophying skeletal muscle: roles and regulation. *Am J Physiol Cell Physiol* 2016; 311: C392-403.
9. Gumucio JP, Mendias CL. Atrogin-1, MuRF-1, and sarcopenia. *Endocrine* 2013; 43: 12-21.
10. Sosa P, Alcalde-Estévez E, Asenjo-Bueno A, Plaza P, Carrillo-López N, Olmos G, *et al.* Aging-related hyperphosphatemia impairs myogenic differentiation and enhances fibrosis in skeletal muscle. *J Cachexia Sarcopenia Muscle* 2021; 12: 1266-1279.

11. Shang GK, Han L, Wang ZH, Liu YP, Yan SB, Sai WW, *et al.* Sarcopenia is attenuated by TRB3 knockout in aging mice via the alleviation of atrophy and fibrosis of skeletal muscles. *J Cachexia Sarcopenia Muscle* 2020; 11: 1104-1120.
12. Xu Y, Tang G, Zhang C, Wang N, Feng Y. Gallic acid and diabetes mellitus: Its association with oxidative stress. *Molecules* 2021; 26: 7115.
13. Bai J, Zhang Y, Tang C, Hou Y, Ai X, Chen X, *et al.* Gallic acid: Pharmacological activities and molecular mechanisms involved in inflammation-related diseases. *Biomed Pharmacother* 2021; 133: 110985.
14. Mori T, Koyama N, Yokoo T, Segawa T, Maeda M, Sawmiller D, *et al.* Gallic acid is a dual α/β -secretase modulator that reverses cognitive impairment and remediates pathology in Alzheimer mice. *J Biol Chem* 2020; 295: 16251-16266.
15. Rahimifard M, Baeeri M, Bahadar H, Moini-Nodeh S, Khalid M, Haghi-Aminjan H, *et al.* Therapeutic effects of gallic acid in regulating senescence and diabetes; an *in vitro* study. *Molecules* 2020; 25: 5875.
16. Guo L, Cao JH, Wei TT, Li JH, Feng YK, Wang LP, *et al.* Gallic acid attenuates thymic involution in the d-galactose induced accelerated aging mice. *Immunobiology* 2020; 225: 151870.
17. Zarei M, Sarihi A, Zamani A, *et al.* Mitochondrial biogenesis and apoptosis as underlying mechanisms involved in the cardioprotective effects of Gallic acid against D-galactose-induced aging. *Mol Biol Rep* 2023; 50: 8005-8014.
18. Zarei M, Sarihi A, Zamani A, Raoufi S, Karimi SA, Ramezani-Aliakbari F. Alginate oligosaccharide alleviates D-galactose-induced cardiac ageing via regulating myocardial mitochondria function and integrity in mice. *J Cell Mol Med* 2021; 25: 7157-7168.
19. Liu X, Liu M. Anti-fatigue effect of ferulic acid in exercise trained mice. *Acta Pol Pharm* 2023; 80: 473-481.
20. Li R, Liu X, Shi Y, Zhou W, Cao L, Li H, *et al.* Application of immunofluorescence in expression detection of Tbx5 in embryonic heart of mice. *J Hunan Univ Arts Sci (Sci Tech)* 2014; 26: 33-36.
21. Priego T, Martín AI, González-Hedström D, Granado M, López-Calderón A. Role of hormones in sarcopenia. *Vitam Horm* 2021; 115: 535-570.
22. Shou J, Chen PJ, Xiao WH. Mechanism of increased risk of insulin resistance in aging skeletal muscle. *Diabetol Metab Syndr* 2020; 12: 14.
23. Asdaq SMB, Alamri AS, Alsanie WF, Alhomrani M, Yasir M, F. Potential benefits of gallic acid as skeletal muscle relevant in animal experimental models. *Saudi J Biol Sci* 2021; 28: 7575-7580.
24. Lee SR, Jo SL, Heo JH, Kim TW, Lee KP, Hong YH. The aqueous fraction of *Castanea crenata* inner shell extract reduces obesity and intramuscular lipid accumulation via induction of mitochondrial respiration and fatty acid oxidation in muscle. *Phytomedicine* 2022; 98: 153974.
25. Wong TS, Mohamed Tap F, Hashim Z, Abdul Majid FA, Zakaria NH, Siahaan P, *et al.* Dual actions of gallic acid and andrographolide trigger AdipoR1 to stimulate insulin secretion in a streptozotocin-induced diabetes rat model. *J Tradit Complement Med* 2022; 13: 11-19.
26. Gandhi GR, Jothi G, Antony PJ, Balakrishna K, Paulraj MG, Ignacimuthu S, *et al.* Gallic acid attenuates high-fat diet fed-streptozotocin-induced insulin resistance via partial agonism of PPAR γ in experimental type 2 diabetic rats and enhances glucose uptake through translocation and activation of GLUT4 in PI3K/p-Akt signaling pathway. *Eur J Pharmacol* 2014; 745: 201-216.
27. Dourado DM, Matias R, da Silva BAK, Milanesi FF, Martello MD, Dos Santos CHM. Benefits of *Sebastiania hispida* (Euphorbiaceae) extract and photobiomodulation therapy as perhaps adjunctive strategies to be explored against snake envenoming. *Photochem Photobiol Sci* 2021; 20: 1069-1085.
28. Gholamine B, Houshmand G, Hosseinzadeh A, Kalantar M, Mehrzadi S, Goudarzi M. Gallic acid ameliorates sodium arsenite-induced renal and hepatic toxicity in rats. *Drug Chem Toxicol* 2021; 44: 341-352.
29. Ojeaburu SI, Oriakhi K. Hepatoprotective, antioxidant and, anti-inflammatory potentials of gallic acid in carbon tetrachloride-induced hepatic damage in Wistar rats. *Toxicol Rep* 2021; 8: 177-185.
30. Yang K, Jian S, Guo D, Wen C, Xin Z, Zhang L, *et al.* Fecal microbiota and metabolomics revealed the effect of long-term consumption of gallic acid on canine lipid metabolism and gut health. *Food Chem X* 2022; 15: 100377.
31. Baharmi S, Kalantari H, Kalantar M, Goudarzi M, Mansouri E, Kalantar H. Pretreatment with gallic acid mitigates cyclophosphamide induced inflammation and oxidative stress in mice. *Curr Mol Pharmacol* 2022; 15: 204-212.
32. Baldin SL, de Pieri Pickler K, de Farias ACS, Bernardo HT, Scussel R, da Costa Pereira B, *et al.* Gallic acid modulates purine metabolism and oxidative stress induced by ethanol exposure in zebrafish brain. *Purinergic Signal* 2022; 18: 307-315.
33. Jiang L, Wu Y, Qu C, Lin Y, Yi X, Gao C, *et al.* Hypouricemic effect of gallic acid, a bioactive compound from *Sonneratia apetala* leaves and branches, on hyperuricemic mice. *Food Funct* 2022; 13: 10275-10290.
34. Wen L, Tang L, Zhang M, Wang C, Li S, Wen Y, *et al.* Gallic acid alleviates visceral pain and depression via inhibition of P2X7 receptor. *Int J Mol Sci* 2022; 23: 6159.
35. Fanaei H, Mard SA, Sarkaki A, Goudarzi G, Khorsandi L. Gallic acid protects the liver against N₂FLD induced by dust exposure and high-fat diet through inhibiting oxidative stress and repressing the inflammatory signaling pathways NF- κ B/TNF- α /IL-6 in Wistar rats. *Avicenna J Phytother* 2021; 11: 527-540.
36. Singla E, Dharwal V, Nigam AS. Gallic acid protects against the COPD-linked lung inflammation and emphysema in mice. *Inflamm Res* 2020; 69: 423-434.
37. Liu X, Liu M. Roles of Ferulic acid on muscle atrophy and grip strength in diabetic mice. *Pakistan Journal of Pharmaceutical Sciences* 2023; 36: 1025-1030.
38. Hong KY, Lee HS, Hong JS, Kim DH, Moon JM, Park Y. Effect of catechinase-converted green tea extract on skeletal muscle development. *BMC Complement Med Ther* 2020; 20: 47.
39. Jia L, Sun S, Ryu Y, Piao ZH, Liu B, Choi SY, *et al.* Gallic acid improves cardiac dysfunction and fibrosis in pressure overload-induced heart failure. *Sci Rep* 2018; 8: 9302.
40. Yigitturk G, Acara AC, Erbas O, Oltulu F, Yavasoglu NUK, Uysal A, *et al.* The antioxidant role of agomelatine and gallic acid on oxidative stress in STZ induced type I diabetic rat testes. *Biomed Pharmacother* 2017; 87: 240-246.
41. Gurkan G, Erdogan MA, Yigitturk G, Erbas O. The restorative effect of gallic acid on the experimental sciatic nerve damage model. *J Korean Neurosurg Soc* 2021; 64: 873-881.
42. Hussein RM, Anwar MM, Farghaly HS, Kandeil MA. Gallic acid and ferulic acid protect the liver from thioacetamide-induced fibrosis in rats via differential expression of miR-21, miR-30 and miR-200 and impact on Tgf- β 1/Smad3 signaling. *Chem Biol Interact* 2020; 324: 109098.
43. Rong Y, Cao B, Liu B, Li W, Chen Y, Chen H, *et al.* A novel Gallic acid derivative attenuates BLM-induced pulmonary fibrosis in mice. *Int Immunopharmacol* 2018; 64: 183-191.
44. Paraíso AE, Sousa JN, Andrade JMO, Mangabeira ES, Lelis DF, de Paula AMB, *et al.* Oral gallic acid improves metabolic profile by modulating SIRT1 expression in obese mice brown adipose tissue: A molecular and bioinformatic approach. *Life Sci* 2019; 237: 116914.
45. Dutta M, Paul G. Gallic acid protects rat liver mitochondria *ex vivo* from bisphenol A induced oxidative stress mediated damages. *Toxicol Rep* 2019; 6: 578-589.
46. Pal C, Bindu S, Dey S, Alam A, Goyal M, Iqbal MS, *et al.* Gallic acid prevents nonsteroidal anti-inflammatory drug-induced gastropathy in rat by blocking oxidative stress and apoptosis. *Free Radic Biol Med* 2010; 49: 258-267.
47. Doan KV, Ko CM, Kinyua AW, Yang DJ, Choi YH, Oh IY, *et al.* Gallic acid regulates body weight and glucose homeostasis through AMPK activation. *Endocrinology* 2015; 156: 157-168.
48. Yan X, Zhang YL, Zhang L, Zou LX, Chen C, Liu Y, *et al.* Gallic

acid suppresses cardiac hypertrophic remodeling and heart failure. *Mol Nutr Food Res* 2019; 63: e1800807.

49. Durgun C, Deveci E. Gallic acid treatment protects intestinal tissue against ischaemia-reperfusion. *Folia Morphol (Warsz)* 2023; 82: 633-640.

50. Jin L, Piao ZH, Liu CP, Sun S, Liu B, Kim GR, *et al.* Gallic acid attenuates calcium calmodulin-dependent kinase II-induced apoptosis in spontaneously hypertensive rats. *J Cell Mol Med* 2018; 22: 1517-1526.

51. Blas-Valdivia V, Franco-Colín M, Rojas-Franco P, Chao-Vazquez A, Cano-Europa E. Gallic acid prevents the oxidative and endoplasmic reticulum stresses in the hippocampus of adult-onset hypothyroid rats. *Front Pharmacol* 2021; 12: 671614.

52. Abdel-Moneim A, Yousef AI, Abd El-Twab SM, Abdel Reheim ES, Ashour MB. Gallic acid and p-coumaric acid attenuate type 2 diabetes-induced neurodegeneration in rats. *Metab Brain Dis* 2017; 32: 1279-1286.

Corrected Proof