

The effects of ATP-dependent potassium channel opener; pinacidil, and blocker; glibenclamide, on the ischemia induced arrhythmia in partial and complete ligation of coronary artery in rats

Selçuk Yaşar¹, Ömer Bozdoğan^{1*}, Salih Tunç Kaya², Hayriye Soytürk Orallar¹

¹ Abant İzzet Baysal University, Faculty of Science and Art, Biology Department 14280 Golkoy, Bolu Turkey

² Duzce University, Faculty of Science and Arts, Biology Department 81620 Konuralp, Duzce Turkey

ARTICLE INFO

Article type:

Original article

Article history:

Received: Mar 9, 2014

Accepted: May 21, 2014

Keywords:

Arrhythmia

Coronary ligation

Ischemia-reperfusion

Myocardial

ABSTRACT

Objective(s): Electrical inhomogeneity between ischemic and non ischemic myocardium is the basis of arrhythmia which occurs following coronary artery occlusion. The leakage of potassium from the ischemic region to the non ischemic region is very effective in the generation of these arrhythmias. The aim of this study is to research the effect of ATP-dependent potassium (K_{ATP}) channel blocker (glibenclamide) and opener (pinacidil) on ischemia induced arrhythmia in the presence of small and large infarct sizes.

Materials and Methods: In this study Sprague-Dawley male rats of 8-9 months of age were used. Ischemia was produced by the partial ligation of left coronary artery ramus descending (PL) for smaller infarct and complete ligation of this artery (CL) for larger infarct for 30 min. The arrhythmia score which was calculated from the duration and type of arrhythmia was significantly higher in animals which had a larger infarct area than the animals which had a smaller infarct.

Results: Glibenclamide increased the rate of arrhythmia in animals having smaller infarct but not in animals having larger infarct. Pinacidil did not affect the occurrence of arrhythmia in either group. There was a significant difference in the infarct size and risk of infarct zone between animals which had small and large infarct sizes. The effect of glibenclamide and pinacidil on the arrhythmias differed depend on decrease of infarct size.

Conclusion: Glibenclamide is not effective to decrease ischemia induced arrhythmia in the presence of small and pinacidil in large ischemic zone.

► Please cite this paper as:

Yaşar S, Bozdoğan Ö, Kaya ST, Soytürk Orallar H. The effects of ATP-dependent potassium channel opener; pinacidil, and blocker; glibenclamide on the ischemia induced arrhythmia change in partial and complete ligation of coronary artery in rats. Iran J Basic Med Sci 2015; 18:188-193.

Introduction

Cardiovascular diseases are the main causes of sudden death in both men and women. About 50% of cardiovascular mortality is caused by coronary artery disease (CAD) (1). The major reason for death related to CAD is the lethal arrhythmias that appear after the coronary occlusion. It is known that one of the factors underlying these arrhythmias is the potassium leakage from the ischemic cells (2-4). The blockage of potassium leakage from ischemic cells can be achieved using an ATP-dependent potassium (K_{ATP}) channel blocker. The antiarrhythmic effects of K_{ATP} channel blockers have been demonstrated in some studies (5-10). But other studies showed that K_{ATP} channel blockers are proarrhythmic in male animals during reperfusion (11, 12). On the other hand, K_{ATP} channel openers have also a proarrhythmic effect (13, 14). The main reason for this

proarrhythmic effect depends on the reentrant arrhythmia caused by the shortening of the action potential duration in myocardial cells (15). However, some researchers reported the antiarrhythmic effect of K_{ATP} channel openers (8, 16 - 18). This different effect may depend on changes in heterogeneity in action potential duration in the ischemic myocardium. This type of heterogeneity is greater between ischemic and non ischemic myocardium. Glibenclamide or pinacidil may decrease the heterogeneity and the likelihood of arrhythmia. The infarct size may be important in the effect of K_{ATP} channel modulators on decreasing electrical inhomogeneity and arrhythmia. The arrhythmia and the rate of death in experimentally produced myocardial infarction have been observed more in animals which have a larger infarct size (19).

Since glibenclamide closes the ischemic myocar-

*Corresponding author: Ömer Bozdoğan. Department of Biology, Faculty of Sciences and Art, Abant İzzet Baysal University, Bolu, Turkey. Tel: +903742541237; Fax: +903742534642; email: bozdogan_o@ibu.edu.tr

dium to the normal myocardium, it may be effective in decreasing electrical inhomogeneity and arrhythmia. There was no study found showing the effect of K_{ATP} channel modulators on arrhythmia in relation to different infarct sizes. In this study, it is hypothesized that glibenclamide decrease electrical inhomogeneity and the occurrence of arrhythmia in the presence of small and pinacidil large infarct sizes.

Materials and Methods

Animals

In this study, male Sprague-Dawley rats 46 (8–9 months old) were used. The animals were allowed to drink tap water *ad libitum* and were fed with commercial rat food pellet. Two main groups were designed; the first group was designated as partial ligation (PL), and the second one was designated as complete ligation (CL). Each main group had three subgroups: control, glibenclamide, pinacidil. All study protocols were approved by the Ethical Committee of Abant İzzet Baysal University, Bolu, Turkey.

Operation

Animals were anesthetized with thiopental sodium (100 mg/kg). The trachea was cannulated for artificial respiration. The catheter for carotid cannulation was filled with heparin (25000 IU/100 ml) diluted 1/10 with saline and inserted into the left carotid artery for the measurement of blood pressure with a blood pressure transducer (SS 13 L, Biopac Systems, Santa Barbara, CA, USA). The chest was opened in the fourth intercostal space. The pericardium was removed, and the heart was exposed.

A loose loop of 5-0 atraumatic silk was quickly placed below the side branch of the left coronary artery ramus descending at the level of the left atrial appendage for small infarct, and around the branching point of the left coronary artery from ramus circumflexus from which only the left arterial descending branch was ligated. Then, the heart was set back in its place and artificial respiration was started using an animal respirator at 60 strokes/min, 9 ml / kg (Ugo Basile Rodent Ventilator, Italy). Subcutaneous needle electrodes were placed under the skin to record the electrocardiogram (ECG). Then, the animals were allowed to stabilize for 5 min. The ligation was made by tightening the silk by producing a bowknot to induce myocardial ischemia for 30 min. Arterial blood pressure and bipolar ECG were recorded over the 30 min of coronary ligation (Biopac Systems, Santa Barbara, CA, USA). At the end of the experiment, live animals were heparinized (500 IU/kg) and the heart was excised. Hearts were perfused with 10 ml of isotonic NaCl solution and then 2 ml of ethanol (96%) for demarcation of the occluded and nonoccluded myocardium (20). After the perfusion, the nonperfused area that remained red was separated from the well perfused area that appeared white in color. The nonperfused

myocardium was cut and weighed. The percentage of this area in respect to the total weight of the ventricle was determined. After that, the non-perfused myocardium was cut into about six segments by razor blade. These segments were put in the solution prepared with nitro blue tetrazolium (NBT) (Sigma-Aldrich CO) about 3 min for staining. The non ischemic region was stained with NBT but the ischemic area was not stained. The color of non ischemic region was light in pallor whereas the infarcted region was red in color.

The infarct region was separated from the noninfarcted region by razor blade and weighed. Finally, the percentage of the risk of arrhythmia to the infarct area (nonperfused area/all of the heart) and infarct area (infarct area/all of the heart) were calculated.

Preparation of drugs

Pinacidil (Sigma-Aldrich CO) was dissolved in ethyl alcohol (96%) and then mixed with isotonic solution and it was given in 0.1 mg/ 0.2 ml / kg doses intravenously from a coccygeal tail vein 10 min before coronary ligation. Glibenclamide (Sigma-Aldrich CO) was dissolved in dimethylsulfoxide and then mixed with ethanol in the same ratio (1/1).

Glibenclamide was given in 5 mg/0.2 ml/kg doses intraperitoneally 20 min before coronary ligation. In the control group, instead of the pinacidil and glibenclamide, the same value of solvent 0.2 ml/kg was applied in both regions. The doses of both drugs used in this study were chosen according to the studies performed previously (21). Since pinacidil was also a hypotensive agent, larger doses usually lead to the dead of animals due to bradycardia and cardiac arrest. Otherwise glibenclamide is more effective in 5 mg/kg dose in intraperitoneal injection. Lower doses of glibenclamide which were tested in previous study were found to be ineffective on ischemia induced arrhythmia (10).

The evaluation of the arrhythmias

The heart rate and blood pressure were determined from recorded ECG before and at 0.5, 1, 10, and 30 min during the 30 min of coronary ligation. The incidence and the total length of arrhythmia were calculated from the recorded ECG. The incidence of arrhythmias was analyzed in accordance with the Lambeth Conventions (22), as ventricular fibrillation (VF), ventricular tachycardia (VT), and other types of arrhythmias including ventricular premature contractions (VPC) and ventricular bigeminy. An arrhythmia score was given according to the incidence and duration of arrhythmias for each animal as follows; 0= No arrhythmias; 1= <10 sec VT or other arrhythmias; 2= 11-30 sec VT or other arrhythmias; 3= 31-90 sec VT or other arrhythmias; 4= 91-180 sec VT or other arrhythmias and/or <10 sec reversible VF; 5=>180 sec VT or other arrhythmias and/or >10 sec reversible

Table 1. Mean heart rate and blood pressure before and following minutes of coronary ligation

	Coronary ligation									
	Heart rate (beats/minute)					Blood pressure (mmHg)				
	Basal	0.5 th min	1 st min	10 th min	30 th min	Basal	0.5 th min	1 st min	10 th min	30 th min
Control PL	369 ± 20	368 ± 20	355 ± 23	346 ± 17	317 ± 11	104 ± 4	91 ± 6	90 ± 8	91 ± 9	74 ± 7
Gli. PL	347 ± 15	352 ± 13	351 ± 18	347 ± 16	354 ± 16*	111 ± 5	86 ± 6	72 ± 9	97 ± 8	98 ± 6*
Pin. PL	350 ± 9	349 ± 9	351 ± 6	348 ± 10	355 ± 7*	119 ± 4*	92 ± 5	93 ± 5	104 ± 6	100 ± 7*
Control CL	367 ± 12	369 ± 12	363 ± 11	379 ± 15	349 ± 14	104 ± 5	70 ± 4*	78 ± 4	92 ± 6	95 ± 7*
Gli. CL	345 ± 12	329 ± 12 ^ε	341 ± 12	328 ± 14 ^ε	323 ± 14	100 ± 6	70 ± 5	68 ± 6	69 ± 6 ^ε	84 ± 7
Pin. CL	390 ± 7	366 ± 7	381 ± 9	364 ± 10	346 ± 8	98 ± 5	61 ± 7	60 ± 8	70 ± 7 ^ε	54 ± 7 ^ε

**P*-value <0.05; according to control PL group. ^ε*P*-value <0.05; according to control CL group. Basal; one min before coronary ligation and at 0.5th, 1st, 10th, and 30th min of coronary ligation, CL; complete ligation of left coronary artery ramus descending, PL; partial ligation of left coronary artery ramus descending, Gli; glibenclamide, Pin; pinacidil. Values represent mean ± SEM

VF; 6= irreversible VF (21).

Statistical analyses

All data are expressed as mean±SEM. The differences between groups were calculated using One-way ANOVA and LSD *post-hoc* test. The survival rate and the incidence of arrhythmia were compared by Chi-square test or Fisher exact test, two tailed.

Results

Changes in the form of QRS complex and/or ST segment elevation were observed in all groups following the coronary artery ligation. All animals survived after 30 min of coronary ligation in the PL group. However, two animals in CL group died due to irreversible fibrillation. Blood pressure significantly decreased immediately after coronary ligation in all groups of animals (*P*-value <0.05) (Table 1). However, the decrease in blood pressure following ligation in CL group was more than the PL group.

Arrhythmias started at 2-5 min following coronary ligation and terminated at about 14 min of coronary ligation. Glibenclamide significantly increased the length of the arrhythmic period in CL group in respect to control (*P*-value <0.05) (Table 2). But it was not effective on the arrhythmic periods of PL group animals.

The length of ventricular tachycardia, total arrhythmia, and other types of arrhythmia were lower in the PL group in respect to the CL group (*P*-value <0.05). Glibenclamide and pinacidil did not

affect the incidence of arrhythmias in either group in respect to the corresponding controls (Table 3).

Arrhythmia started earlier in the PL group than the CL group. Pinacidil lengthened the onset of arrhythmia in the PL group but not in the CL group. The effect of both glibenclamide and pinacidil on arrhythmic period was found to be similar in the PL and CL groups. Glibenclamide increased the arrhythmic period significantly in CL group (*P*-value <0.05) but non significantly in PL group (*P*-value = 0.61), but pinacidil non significantly decreased the arrhythmic period in the CL (*P*-value = 0.46) and PL (*P*-value = 0.35) groups. As expected, the score of arrhythmia was significantly higher in the CL group than the PL group (*P*-value <0.05). Glibenclamide significantly increased the score of arrhythmia in PL group (*P*-value <0.05) but the effects of pinacidil were non significant (*P*-value = 0.21) (Figure 1). Both glibenclamide and pinacidil did not affect the score of arrhythmia in CL group.

The risk of infarct zone and infarct area was larger in the CL group than the PL group (*P*-value <0.05). Glibenclamide and pinacidil non significantly increased the infarct area in the PL group (*P*-value = 0.24) but they were not effective in the CL group, (Table 4).

Risk of infarct area% = weight of risk of infarct zone × 100/weight of whole ventricular myocardium.

Infarct area% = weight of infarct area × 100/weight of whole ventricular myocardium. **P*-value < 0.05; according to control CL group.

Table 2. Duration of arrhythmias occurred in 30 minute coronary ligation


Groups	N	Onset of arrhythmia (min)	Arrhythmic period (min)	Length of arrhythmia (min)			
				VF	VT	Other	Total
Control PL	7	2.18 ± 0.94*	13.09 ± 3.40	0	0.023 ± 0.015*	0.44 ± 0.18*	0.46 ± 0.18*
Gli. PL	7	4.40 ± 1.04	14.61 ± 1.18	0.026 ± 0.017	0.03 ± 0.02	1.1 ± 0.25	1.15 ± 0.26
Pin. PL	7	4.88 ± 0.91 ^α	10.30 ± 1.21	0	0	0.8 ± 0.14	0.79 ± 0.14
Control CL	8	5.88 ± 0.57	15.44 ± 2.22	0	1.02 ± 0.44	1.68 ± 0.28	2.7 ± 0.71
Gli. CL	7	3.99 ± 0.94	21.57 ± 1.64*	0.021 ± 0.017	0.48 ± 0.47	2.64 ± 0.68	3.15 ± 0.79
Pin. CL	8	5.82 ± 0.26	13.38 ± 1.71	0.06 ± 0.06	0.54 ± 0.48	2.08 ± 0.38	2.68 ± 0.85

**P*-value <0.05; according to control CL group, ^α*P*-value <0.05; according to control PL group, N: the number of animals at the end of 30 min of ligation, VF: ventricular fibrillation, VT: ventricular tachycardia, Other: ventricular extra beats and bigeminy CL: complete ligation of left coronary artery ramus descending, PL: partial ligation of left coronary artery ramus descending Gli: glibenclamide, Pin: pinacidil. Values represent mean ± SEM

Table 3. The incidence of arrhythmias during 30 minute coronary ligation in male rats

Group	N	Survived (n/%)	Incidence of arrhythmias (n/%)			
			VF	VT	Other	Bradycardia
Control PL	7	7/100	0/0	2/29	7/100	0/0
Gli. PL	7	7/100	2/29	2/29	7/100	0/0
Pin. PL	7	7/100	0/0	0/0	7/100	0/0
Control CL	10	8/80	2/20	4/40	10/100	0/0
Gli. CL	7	7/100	2/29	2/29	7/100	0/0
Pin. CL	8	8/100	1/13	4/50	8/100	0/0

N: number of animals at the beginning of the ligation, (n/%) : number of animals/percentage of animals in respect to all animals in the group VF: ventricular fibrillation, VT: ventricular tachycardia, Other: ventricular extra beats and bigeminy, CL: complete ligation of left coronary artery ramus descending, PL: partial ligation of left coronary artery ramus descending. Gli: glibenclamide, Pin: pinacidil

**Figure 1.** The score of arrhythmia. *P-value <0.05; according to CL control Group, α P-value <0.05; according to PL control group

Discussion

No study is found to show the effect of K_{ATP} channel on arrhythmias in animals with both small and large infarcts. But it was shown that a gradually increased infarct size increases the severity of arrhythmia (19). This was supported by the results of the present study.

The occurrence of arrhythmia increased in animals with larger infarct size. The leakage of potassium ion from the non ischemic region to the normal myocardium is responsible in the generation of these inhomogeneities and the resulting arrhythmias (2-4). In a previous recent study (23) and earlier study performed by other scientists (8), it was suggested that K_{ATP} channel blockage decreases electrical inhomogeneity in ischemic and non ischemic regions and decreases the probability of the occurrence of arrhythmia. K_{ATP} channel blockage increases the duration of action potential in ischemic cells but not in non ischemic cells (24, 25). It is expected that sarcolemmal K_{ATP} channel blockers block the potassium outflow from ischemic cells to non ischemic region and it closes the ischemic cell to non ischemic cell. By this mechanism the glibenclamide was thought to decrease the arrhythmia in animals with especially small infarct size. But it did not decrease the arrhythmia, instead oppositely increased the arrhythmia. This opposite effect of glibenclamide might be due to its ischemic effect. Animals with smaller infarct size have less number of ischemic cells, so the blockage of the channel might be more plausible and thought to be more effective to decrease arrhythmia. In this case, conversion of ischemic cell to non ischemic cell by

the effect of glibenclamide was thought to decrease electrical inhomogeneity. But this hypothesis was not supported by the result of this study, because the ischemic effect of glibenclamide prevailed instead of decreasing electrical inhomogeneity. Otherwise, this study showed that the effect of glibenclamide is changed depend on the risk of infarct zone, Pinacidil opens the channel in non ischemic cells, decreases action potential duration, and may close the non ischemic cell to ischemic cell thus decreasing the electrical inhomogeneity. In addition, the contractility of myocardium decreases through the effect of pinacidil and leads to the decrease of myocardial oxygen consumption, infarct size (26), and the rate of occurrence of arrhythmia (21) following coronary occlusion. Although glibenclamide closes the ischemic cell to the non ischemic cell, it increases myocardial contraction and infarct production during myocardial ischemia. The proarrhythmic effect of glibenclamide shown in some studies (11, 12) might be attributable to this effect. Pinacidil has opposite effect of glibenclamide at cellular level. It decreases action potential duration and is effective to the normal myocardial cells but not to the ischemic cells. This is the reason that pinacidil was thought to close the non ischemic cell to ischemic cell and again would decrease electrical inhomogeneity in animals have larger risk of infarct zone. Since the animal with larger infarct size has smaller non ischemic cell, the conversion of non ischemic cell to ischemic cell was thought to be more plausible and pinacidil would significantly be effective to decrease arrhythmia in this group. But pinacidil was not found to be effective on the arrhythmia in animals having larger infarct size.

Table 4. The percentage of risk of infarct zone and infarct area measured at the end of 30 minutes coronary ligation

Groups	N	Risk of infarct zone %	Infarct area %
Control PL	7	29.42 ± 1.87*	7.85 ± 2.20*
Gli. PL	7	33.00 ± 2.00	10.28 ± 1.71
Pin. PL	7	30.42 ± 2.18	10.28 ± 0.99
Control CL	10	48.01 ± 1.45	20.99 ± 1.41
Gli. CL	7	45.87 ± 1.16	20.70 ± 1.13
Pin. CL	8	51.68 ± 1.52	22.08 ± 0.65

*P-value <0.05; according to control CL group. N: the number of animals at the end of 30 min of ligation CL: complete ligation of left coronary artery ramus descending, PL: partial ligation of left coronary artery ramus descending, Gli: glibenclamide, Pin: pinacidil. Values represent mean ± SEM

Glibenclamide did not affect the arrhythmia following complete ligation which was similar with the findings of our previous study (27). In addition, pinacidil did not have significant effect on arrhythmias in either animal with small or large infarcts produced by 30 min of ligation. In contrast to the most of the studies, pinacidil did not have proarrhythmic (8, 17) or antiarrhythmic effects (13) following ligation. However, it was seen that pinacidil increased the severity of arrhythmia non significantly in animals with smaller infarcts size.

Glibenclamide and pinacidil increased the infarct area non significantly in animal having partial coronary ligation, but they did not have any effect on the ischemia induced arrhythmia in animals which had complete ligation. Arrhythmia started earlier in group with smaller infarct size that may be caused by lower reflex activation of sympathetic system. The fall of the blood pressure immediately following ligation was lower in the PL group than the CL group which may be attributable to the short lasting sympathetic overdrive suppression in the PL group. Pinacidil did not have any effect on the arrhythmic period in either group of animals. Since no comparable study was found in the literature, the results related with the arrhythmic period could not be compared.

Conclusion

As a result, it seems that the effect of glibenclamide and pinacidil on the severity of arrhythmia changes depend on the change of the ischemic zone size. Glibenclamide increased arrhythmia in the presence of smaller risk of infarct zone but was ineffective in the presence of larger infarct size. Pinacidil did not change the severity of arrhythmia depend on the changes in ischemic zone, but it non significantly increased the total arrhythmia in the presence of smaller ischemic zone. Underlying cellular mechanisms of different effect of glibenclamide and pinacidil on the ischemia induced arrhythmia in small and large infarct size are not explained by the result of present study. Further electrophysiologic researches are required in this case. Since the electrical inhomogeneity is important in the generation of ischemia induced arrhythmias, more drugs or procedures should be researched to decrease electrical inhomogeneity.

Acknowledgment

This study was supported financially by the research fund of Abant İzzet Baysal University, Bolu Turkey. Some part of the results described in this paper is presented as MSc thesis in Biology Department of Faculty of Science, Abant İzzet Baysal University.

References

1. Pokorny J, Stanek V, Vrana M, Sudden Cardiac Death Thirty Years Ago and at present. The role of Autonomic

disterbunces in acute myocardial infarction revisited. *Physiol. Res* 2011; 715-728.

2. Harris AS. Potassium and experimental coronary occlusion. *Am Heart J* 1966; 71: 797-802.

3. Coronel R, Fiolet JW, Wilms-Schopman FJG, Schaapherder AF, Johnson TA, Gettes LS, *et al*. Distribution of extracellular potassium and its relation to electrophysiological changes during acute myocardial ischemia in the isolated perfused porcine heart. *Circulation* 1988; 77: 1125-1138.

4. Wilde AAM, Escande D, Schumacher CA, Thuringen D, Mestre M, Fiolet JWT, *et al*. Potassium accumulation in the globally ischemic mammalian heart. A role for the ATP sensitive potassium channel. *Circ Res* 1990; 67: 835-843.

5. Billman GE, Avendora CE, Halliwill JR. The effects of ATP-dependent potassium channel antagonist, glyburide, on coronary flow and susceptibility to ventricular fibrillation. *J Cardiovasc Pharm* 1993; 21: 197-207.

6. Colatsky TJ, Argentieri TM. Potassium channel blockers as antiarrhythmic drugs. *Drug Develop Res* 1994; 33: 235-249.

7. Tosaki A, Hellegouarch A. Adenosine triphosphate sensitive potassium channel blocking agent ameliorates, but the opening agent aggravates, ischemia/reperfusion induced injury. *JACC* 1994; 23: 487-496.

8. Lepran I, Baczkó I, Varro A, Papp JGy. ATP sensitive potassium channel modulators: Both pinacidil and glibenclamide produce antiarrhythmic activity during acute myocardial infarction in conscious rats. *J Pharmacol Exp Ther*. 1996; 277: 1215-1220.

9. Bozdoğan O. ATP-dependent potassium channel openers and blockers: proarrhythmic or antiarrhythmic effects following myocardial ischemia and reperfusion. *Kafkas Univ Vet Fak Derg* 1997; 3: 127-133.

10. El-Reyani N, Bozdoğan O, Baczkó I. Comparison of the efficacy of glibenclamide and glimepride in reperfusion induced arrhythmias in rats. *Eur J Pharmacol* 1999; 192: 365-187.

11. Bernauer W. Concerning the effect of the K_{ATP} channel blocking agent glibenclamide on ischemic and reperfusion arrhythmias. *Eur J Pharmacol* 1997; 326: 147-156.

12. Del Valle HF, Lascano EC, Negroni JA, Crottogini AJ. Glibenclamide effects on reperfusion-induced malignant arrhythmias and left ventricular mechanical recovery from stunning in conscious sheep. *Cardiovasc Res* 2001; 50: 474-485.

13. Chi L, Uprichard AC, Lucchesi BR. Profibrillatory actions of pinacidil in a conscious canine model of sudden coronary death. *J Cardiovasc Pharm* 1990; 15: 452-464.

14. Tosaki A, Szerdahelyi P, Engelman RM. Potassium channel openers and blockers: Do they proarrhythmic or antiarrhythmic activity in ischemic and reperfused rat hearts? *J Pharmacol Exp Ther* 1993; 263: 1261-1268.

15. Wilde AAM. KATP channels opening and arrhythmogenesis. *J Cardiovasc Pharm* 1994; 24: 35-40.

16. Kerr MJ, Wilson R, Shangs RG. Suppression of ventricular arrhythmias after coronary artery ligation by pinacidil, a vasodilator drug. *J Cardiovasc Pharm* 1985; 7: 875-883.

17. Baczko I, Lepran I, Papp JGy. K_{ATP} channel modulators increase survival rate during coronary occlusion-reperfusion in anaesthetized rats. *Eur J Pharmacol* 1997; 324: 77-83.
18. Bozdogan O, Gonca E, Nebigil M, Tiryaki ES. Effect of thimerosal on arrhythmia induced by coronary ligation. *Int Heart J* 2005; 46: 711-721.
19. Bozdogan O, Bolukbası F. The arrhythmias occurring in the late period of experimentally induced myocardial infarction in dogs. *Turk J Vet Anim Sci* 1994; 8: 147-151.
20. Lepran I, Koltai M, Siegmund W, Skekeres L. Coronary artery ligation, early arrhythmias and determination of the ischemic area in conscious rats. *J Pharmacol Method* 1983; 9: 219-230.
21. Gonca E, Bozdogan O. Both mitochondrial K_{ATP} channel opening and sarcolemmal K_{ATP} channel blockage confer protection against ischemia/-reperfusion- induced arrhythmia in anesthetized male rats. *J Cardiovasc Pharm* 2010; 15: 403-411.
22. Walker MJ, Curtis MJ, Hearse DJ, Campbell RW, Janse MJ, Yellon DM, *et al.* The Lambeth Conventions: guidelines for the study of arrhythmias in ischaemia infarction, and reperfusion. *Cardiovasc Res* 1988 Jul; 22:447-55.
23. Gonca E, Tiryaki SE, Bozdogan O. The effect of sex on the ischemia- reperfusion arrhythmias and the role of ATP-dependent potassium channel blockage. *Turk J Biol* 2004; 28: 39-46.
24. McKensie I, Saville VL, Waterfall JF. Differential class III and glibenclamide effects on action potential duration in guinapig papillary muscle during normoxia and hypoxia/ischemia. *Brit J Pharmacol* 1993; 110: 531-583.
25. Tweedie D, Henderson C, Kane K. Glibenclamide, but not class 3 drugs, prevents ischemic shortening of the refractory period in guinea-pig hearts. *Eur J Pharmacol* 1993; 240: 251-257.
26. Escande D, Caverio L. K⁺ channel openers and natural cardioprotection. *Trends Pharmacol Sci* 1992; 13: 269-272.
27. Bozdogan O, Lepran I, Papp JGy. Effect of the combination of glibenclamide, an ATP-dependent potassium channel blocker, and metoprolol, a cardioselective β -adrenoceptor blocker, during myocardial infarction in conscious rats. *Turk J Med Sci* 2000; 30: 517-522.