

Mangosteen peel extract (*Garcinia mangostana* L.) as protective agent in glucose-induced mesangial cell as *in vitro* model of diabetic glomerulosclerosis

Wahyu Widowati^{1*}, Dian Ratih Laksmiawati², Teresa Liliana Wargasetia¹, Ervi Afifah³, Annisa Amalia³, Yukko Arinta³, Rizal Rizal³, Tri Suciati⁴

¹ Faculty of Medicine, Maranatha Christian University, Bandung 40164, West Java, Indonesia

² Faculty of Pharmacy, Pancasila University, Jakarta Selatan 12630, DKI Jakarta, Indonesia

³ Aretha Medika Utama, Biomolecular and Biomedical Research Center, Bandung 40163, West Java, Indonesia

⁴ School of Pharmacy Bandung Institute of Technology, Bandung 40132, West Java, Indonesia

ARTICLE INFO

Article type:

Short communication

Article history:

Received: Jan 23, 2018

Accepted: Mar 18, 2018

Keywords:

Fibronectin
Glomerulosclerosis
Garcinia mangostana
Mesangial cell
Transforming growth factor- β 1

ABSTRACT

Objective(s): This study aims to evaluate the activity of mangosteen peels extract (MPE) as protection agent on induced-glucose mesangial cells (SV40 MES 13 cell line (*Glomerular Mesangial Kidney, Mus Musculus*)).

Materials and Methods: MPE was performed based on maceration method. Cytotoxic assay was performed based on MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) method, while the level of TGF- β 1 (Transforming growth factor- β 1) and fibronectin in glucose-induced mesangial cells were assayed and determined using ELISA KIT.

Results: In viability assay, MPE 5 and 20 μ g/ml has the highest activity to increase cells proliferation in glucose-induced mesangial cells at 5, 10, and 15 days of incubation in glucose concentration (5 and 25 mM) ($P < 0.05$). In inhibitory activity of TGF- β 1 and fibronectin level, MPE 5 μ g/ml (glucose-induced 5 mM) show the lowest level compared to positive control and other treatments ($P < 0.05$).

Conclusion: MPE can increase cell proliferation in glucose-induced mesangial cells and significantly reduce the level of TGF- β 1 and fibronectin. MPE activity has correlates to inhibit the diabetic glomerulosclerosis condition and may increase mesangial cell proliferation.

► Please cite this article as:

Widowati W, Laksmiawati DR, Wargasetia TL, Afifah E, Amalia A, Arinta Y, Rizal R, Suciati T. Mangosteen peel extract (*Garcinia mangostana* L.) as protection agent in glucose-induced mesangial cell as *in vitro* model diabetic glomerulosclerosis. Iran J Basic Med Sci 2018; 21:972-977. doi: 10.22038/IJBMS.2018.29349.7094

Introduction

According to the International Diabetes Federation, the prevalence of diabetes in the world is estimated to increase from 285 million persons to 439 million in 2030 (1). Diabetic is known to be a leading cause of end-stage renal failure (2). All forms of diabetes are characterized by hyperglycemia (3). Hyperglycemia is the primary pathogenic factor for diabetic nephropathy (DN) (4). Through multiple mechanisms, diabetic nephropathy can develop to end-stage kidney disease but none is as important as the gradual, inexorable scarring of the renal glomerulus, known as glomerulosclerosis (5). Glomerulosclerosis is diabetic nephropathy caused by accumulation of extracellular matrix (ECM) proteins in mesangial interstitial space, resulting in fibrosis manifested by either diffuse or nodular changes (6). One of the most common matrix protein detected is fibronectin (5). Several studies also found that hyperglycemia induces reactive oxygen species (ROS) production in mesangial cells that up-regulates Transforming Growth Factor Beta (TGF- β) involved in ECM accumulation (7, 4).

During the centuries, natural substances from plant has been widely used for treating and preventing some various diseases. Most of these natural substances were studied, isolated, and converted into modern medicine

(8). These natural substances and its compounds would be promising alternative for therapeutic in respect of low cost, highly compatible with dietary intake and no harmful effects inside the human body (9). *Garcinia mangostana* Linn. or commonly known as mangosteen is a tropical fruit from South East Asia (10). Not only known from its flesh as a dessert, the peels of mangosteen are also known to treat various infectious diseases of skin and wounds, diarrhea, dysentery, cholera and have anti-inflammatory (11-13), anticancer potency (14). The peels of mangosteen are reported to be rich of phenolic compounds with potential applications as therapeutic agents such as phenolic acids (15), tannins (16), xanthenes and anthocyanins (17, 10, 13). From these various activities, the potential of mangosteen peels against atherosclerosis is thought to be derived from antioxidant (18), antiaggregation (19), antiobesity (20, 21, 22) anti-inflammatory properties (13). Mangosteen peel extract (MPE) containing many active compounds are expected to inhibit and retard progression of diabetic glomerulosclerosis into renal chronic disease. This research was conducted to evaluate the potential of MPE and its component α -mangostin (AM) as protective agent in glucose-induced mesangial cell as *in vitro* model diabetic glomerulosclerosis.

*Corresponding author: Wahyu Widowati. Faculty of Medicine, Maranatha Christian University, Bandung 40163, West Java, Indonesia. Tel/ Fax: +62-81910040010; Email: wahyu_w60@yahoo.com

Materials and Methods

Plants extract preparation

G. mangostana L. was collected from Cisalak-Subang, West Java, Indonesia plantation and identified by a staff of Herbarium of Department of Biology, School of Life Science and Technology, Bandung Institute of Technology, Bandung, West Java, Indonesia. The mangosteen peels were collected, chopped, and kept in drier tunnel service. Extraction was performed based on maceration method using distilled ethanol 70% as the solvent for collecting *G. mangostana* L. peel extract (MPE) (13,19).

Viability assay

The Glomerular Mesangial Kidney, *Mus musculus* (SV40 MES 13 ATCC ® CRL-1927™) was obtained from Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung. The viability assay was performed using MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) Proliferation Assay Kit (Abcam, ab197010). In brief, 5×10^3 cells per well in F12-K medium (Gibco, 21127022) and DMEM (Gibco, 11995065), 10% fetal bovine serum (FBS, Gibco, 10270106), and 1% antibiotic-antimycotic (Gibco, 1772653), 1% HEPES (Sigma Aldrich, 1002184736) were cultured in 96 well plate (Corning, 3596) and incubated at 37 °C, 5% CO₂ for 24 hr. Then, medium replaced with 180 µl of fresh medium, 20 µl of MPE (5 µg/ml and 20 µg/ml), AM (20 µM and 80 µM) and DMSO 10% were added in triplicate and the plates were incubated at 37 °C, 5% CO₂ for 24 hr. Untreated cells were served as the control. Briefly, 20 µl MTS was added to each well. Then, the plate was incubated at 37 °C, 5% CO₂ for 4 hr. The absorbance was measured at 490 nm using Multiskan GO plate reader (Thermo Scientific, U.S.A) (23, 24, 13).

Glucose-induced mesangial cells for proliferation assay

Briefly 5×10^3 cells/well of SV40 MES 13 cells were plated in 96-well plate with 200 µl growth medium and incubated at 37 °C, 5% CO₂ for 24 hr. The medium was discarded then added with 180 µl glucose-induced medium (5 mM, 20 mM, 50 mM, and 115 mM) 20 µl MPE (5, 20 µg/ml) and 20 µl AM (20, 80 µM). After that, the cells were incubated at 37 °C, 5% CO₂ for 14 days. Proliferation was measured every 2 days using MTS Proliferation Assay Kit (Abcam, ab197010). The absorbance was measured at 490 nm using Multiskan GO plate reader (Thermo Scientific, U.S.A) to calculate the percentage of cell mortality (25, 13).

Quantification of TGF-β1 level

The quantitative determination of TGF-β1 level in the cell-free supernatant was performed using Rat TGF-β1 ELISA Kit (ElabSci E-EL-R0084) based on manufactured protocol. Briefly, 100 µl of standard, blank, and sample solution was added into each well then sealed and incubated for 90 min at 37 °C. After treating with MPE and AM, the cell-free supernatant was served as the sample. The glucose-induced mesangial cell free supernatant without extract and compounds were used as positive control. The normal cell or untreated cell was

used as negative control. Subsequently, the liquid of each well was discarded and 100 µl biotinylated detection Ab was added and then incubated for an hr at 37 °C. Then the liquid was discarded and the plate was washed three times using 200 µl wash buffer. HRP conjugate (100 µl) was added and incubated for 30 min at 37 °C. The liquid was discarded again and the plate was washed five times using 200 µl wash buffer. Substrate reagent (90 µl) was added and incubated for 15 min at 37 °C. Stop solution (50 µl) was added and the absorbance was read at 450 nm using Multiskan GO plate reader (Thermo Scientific, U.S.A) (26).

Quantification of fibronectin level

The quantitative determination of fibronectin level in the cell-free supernatant was performed using Rat FN (Fibronectin) ELISA Kit (ElabSci E-EL-R0578) based on manufactured protocol. Briefly, 100 µl of standard, blank, and sample solution was added into each well then sealed and incubated for 90 min at 37 °C. The cell-free supernatant, after treated with MPE and AM, were served as the sample. The glucose-induced mesangial cell free supernatant without extract and compounds were used as positive control. The normal cell or untreated cell was used as negative control. Subsequently, the liquid of each well was discarded and 100 µl of biotinylated detection Ab was added then incubated for an hour at 37 °C. The liquid was discarded again and the plate was washed three times using 200 µl wash buffer. HRP conjugate (100 µl) was added and incubated for 30 min at 37 °C. The liquid was discarded again and the plate was washed five times using 200 µl wash buffer. Substrate reagent (90 µl) was added and incubated for 15 min at 37 °C. Stop solution (50 µl) was added and the absorbance was measured at 450 nm using Multiskan GO plate reader (Thermo Scientific, U.S.A) (27).

Statistical analysis

The data was analyzed using SPSS 16 (SPSS Inc., Chicago, IL, USA) to perform one-way ANOVA to verify the results of different treatments and Duncan *post hoc* was used to validate significant differences for all treatments ($P < 0.05$). The results are displayed as means ± standard deviation.

Results

Viability assay

The viability of glucose-induced SV40 MES 13 cells during 5 days of incubation time treated with MPE (5, 20 µg/ml) and AM (20, 80 µM) can be seen in Table 1. Cell treated with MPE 5 µg/ml showed the highest cell proliferation. This indicated that MPE 5 µg/ml has a good viability and activity to increase cell proliferation in glucose-induced mesangial cells at 5 days of incubation compared to other treatments.

The viability of glucose-induced SV40 MES 13 cells treated with MPE (5, 20 µg/ml) and AM (20, 80 µM) during 10 days of incubation time can be seen in Table 2. MPE 20 µg/ml has the highest viability cell in all glucose concentration compared to control and other treatments. This data indicated that MPE 20 µg/ml has the highest activity to increase cells proliferation in

Table 1. Effect of mangosteen peel extract (PEM) and α -mangostin (AM) toward cells proliferation in glucose-induced mesangial cells at 5 days of incubation

Treatment	Glucose Concentration			
	0 mM	5 mM	25 mM	125 mM
Control	100.00 \pm 5.84 ^a	100.00 \pm 5.84 ^a	100.00 \pm 5.84 ^a	100.00 \pm 5.84 ^{bc}
Positive Control		96.71 \pm 0.91 ^a	84.78 \pm 1.71 ^a	74.92 \pm 1.30 ^a
MPE 20 μ g/ml	136.32 \pm 28.75 ^{ab}	207.70 \pm 7.02 ^b	162.18 \pm 8.56 ^{bc}	121.04 \pm 12.52 ^c
MPE 5 μ g/ml	153.94 \pm 26.71 ^b	214.61 \pm 3.19 ^b	199.38 \pm 19.80 ^c	121.20 \pm 12.43 ^c
AM 80 mM	113.39 \pm 26.71 ^{ab}	175.18 \pm 42.90 ^b	153.46 \pm 5.35 ^c	99.28 \pm 2.04 ^{bc}
AM 20 mM	143.33 \pm 3.17 ^{ab}	159.76 \pm 25.46 ^b	193.13 \pm 24.44 ^b	98.30 \pm 7.75 ^b

*Data were served in average \pm standard deviation. Different superscript letters in the same column of 0 mM (a, ab, b), 5 mM (a, b), 25 mM (a, b, c), 125 mM (a, b, bc, c) glucose concentration show significant differences among treatments per induction ($P < 0.05$) analyzed using ANOVA and Duncan *post hoc* test

Table 2. Effect of mangosteen peel extract and α -mangostin toward cells proliferation in glucose-induced mesangial cells at 10 days of incubation

Treatment	Glucose Concentration			
	0 mM	5 mM	25 mM	125 mM
Control	100.00 \pm 8.57 ^a	100.00 \pm 8.57 ^{ab}	100.00 \pm 8.57 ^{bc}	100.00 \pm 8.57 ^c
Positive Control		94.64 \pm 0.43 ^a	86.84 \pm 1.50 ^a	73.26 \pm 2.25 ^a
MPE 20 μ g/ml	199.39 \pm 20.50 ^b	127.68 \pm 3.33 ^d	122.02 \pm 3.10 ^d	91.93 \pm 0.15 ^{bc}
MPE 5 μ g/ml	198.59 \pm 15.13 ^b	110.91 \pm 6.34 ^{bc}	111.26 \pm 5.52 ^{cd}	89.79 \pm 0.22 ^b
AM 80 mM	145.65 \pm 8.69 ^{ab}	115.64 \pm 0.64 ^{cd}	101.14 \pm 0.53 ^{bc}	91.88 \pm 0.34 ^{bc}
AM 20 mM	174.85 \pm 40.73 ^b	101.15 \pm 0.27 ^{ab}	98.53 \pm 1.61 ^{ab}	91.49 \pm 1.61 ^{bc}


*Data were served in average \pm standard deviation. Different superscript letters in the same column of 0 mM (a, ab, b), 5 mM (a, ab, bc, cd, d), 25 mM (a, ab, bc, cd, d), 125 mM (a, b, bc, c) glucose concentration show significant differences among treatments per induction ($P < 0.05$) analyzed using ANOVA and Duncan *post hoc* test. Mangosteen peel extract (MPE)

glucose-induced mesangial cells at 10 days of incubation.

The viability of glucose-induced SV40 MES 13 cells treated with MPE (5, 20 μ g/ml) and AM (20, 80 μ M) during 15 days of incubation time can be seen in Table 3. In each glucose concentration, the highest cell proliferation was MPE 20 μ g/ml in glucose concentration 5 and 25 mM with each value of 113.38 \pm 5.10% and 97.85 \pm 0.58%. This indicated that MPE 20 μ g/ml in 5 and 25 mM glucose concentration is potential to increase cell proliferation in glucose-induced mesangial cells at 15 days of incubation. Based on statistical analysis, MPE has significant difference in elevation of cell proliferation in glucose-induced mesangial cells compared to positive control ($P < 0.05$).

TGF- β 1 level

The reduction of TGF- β 1 level in glucose-induced mesangial cells as positive control, glucose-induced mesangial cells treated with MPE (5 and 20 μ g/ml) and AM (20 and 80 μ M) can be seen in Figure 1. Figure 1 showed the concentration of TGF- β 1 by ELISA method after treating glucose-induced SV40 MES 13 cells with MPE and AM. The lowest level of TGF- β 1 was obtained in 5 μ g/ml of MPE (66.30 pg/ml) at 5 mM glucose-induced, while AM was at 80 μ M (65.42 pg/ml) at 5 mM glucose-

**Figure 1.** TGF- β 1 level of glucose-induced SV40 MES 13 cells treated with mangosteen peel extract and α -mangostin

*Data were analyzed with ANOVA and Duncan *post hoc* test. The different letter (a, ab, b, bc, cd, de, e, f) show significant differences among treatments ($P < 0.05$). Each samples were done in triplicate. Negative control was un-induced mesangial cells, positive control I was glucose 5 mM-induced cells, positive control II was glucose 10 mM-induced cells

induced compared to positive control (5 mM glucose-induced) (95.58 pg/ml). Based on statistical analysis, MPE 5 μ g/ml has the most significant difference in

Table 3. Effect of mangosteen peel extract and α -mangostin toward cells proliferation in glucose-induced mesangial cells at 15 days of incubation

Treatment	Glucose Concentration			
	0 mM	5 mM	25 mM	125 mM
Control	100.00±7.75	100.00±7.75 ^{ab}	100.00±7.75 ^b	100.00±7.75 ^b
Positive Control		76.89±2.13 ^a	70.94±1.80 ^a	54.12±4.18 ^a
MPE 20 μ g/ml	116.16±4.04	113.38±5.10 ^b	97.85±0.58 ^b	59.11±13.86 ^a
MPE 5 μ g/ml	118.12±3.90	104.60±15.85 ^{ab}	83.32±2.17 ^{ab}	68.42±8.11 ^a
AM 80 mM	122.34±12.65	100.17±11.03 ^{ab}	78.08±4.61 ^a	62.08±8.75 ^a
AM 20 mM	121.98±16.65	104.83±13.32 ^{ab}	72.85±11.71 ^a	70.31±5.27 ^a

*Data were served in average±standard deviation. Different superscript letters in the same column of 5 mM (a, ab, b), 25 mM (a, ab, b), 125 mM (a, b) glucose concentration show significant differences among treatments per induction ($P<0.05$) analyzed using ANOVA and Duncan *post hoc* test




Figure 2. Fibronectin level of glucose-induced SV40 MES 13 cells treated with mangosteen peel extract and α -mangostin.

*Data were analyzed with ANOVA and Duncan *post hoc* test. The different letters (a, ab, bc, c, d, e) show significant differences among treatments ($P<0.05$). Each samples were done in triplicate. Negative control was un-induced mesangial cells, positive control I was glucose 5 mM-induced cells, positive II was glucose 10 mM-induced cells

inhibition of TGF- β 1 level of glucose-induced mesangial cells compared to positive control but almost comparable with negative control ($P<0.05$).

Fibronectin level

The reduction of fibronectin level in glucose-induced mesangial cells as positive control, glucose-induced mesangial cells treated with MPE (5 and 20 μ g/ml) and AM (20 and 80 μ M) can be seen in Figure 2. Figure 2 showed the concentration of fibronectin level by ELISA method after treating glucose-induced SV40 MES 13 cells with MPE and AM. The lowest fibronectin level was obtained at 5 μ g/ml of MPE (1.11 ng/ml) at 5 mM glucose-induced and 80 μ M of AM (2.15 ng/ml) at 5 mM glucose-induced compared to positive controls (5 mM glucose-induced) (7.34 ng/ml). Based on statistical analysis, MPE 5 and 20 μ g/ml has the most significant difference compared to positive control but almost comparable compared to negative control ($P<0.05$).

Discussion

Hyperglycemia is believed to play a pivotal role for the initiation of pathological process. The primary injury is

believed to take place in the glomerular tuft and leads to an eventual decline in renal function (28). According to Qian *et al.* (2014), excessive amount of extracellular glucose leads to glucose uptake in mesangial cells which further leads to an activation of a number of metabolic pathways that results in increased production of reactive oxygen species (ROS) (29) and advanced glycation end products (AGEs). Thus pathways induce ECM production such as fibronectin and critically TGF- β 1 synthesis. Yet, overexpression of TGF- β and fibronectin closely linked to glomerulosclerosis (30).

In this study, mesangial cells were cultured in a high concentration of glucose (hyperglycemic) which correlates to diabetic glomerulosclerosis condition. TGF- β and fibronectin was used as a parameter due to their existence in glomerulosclerosis disease. In line with previous study conducted by Nahman *et al.* (1992), there were significant decreasing of cell number after 5, 10 and 15 days incubation in high concentration of glucose (5, 25, and 125 mM). The results suggest that high concentration of glucose may have suppressive effect on mesangial cell proliferation. In addition, the results showed the effect of glucose on cell proliferation is dose dependent (25). The longer the incubation and higher the glucose concentration, the ability of the cells to improve itself is lower. In other study, it also suggested that a high glucose in mesangial cell reduced cell number caused by free radical damage and enhanced ECM (31). Yet, the mechanisms of how glucose inhibit cell proliferation is remain unclear but the production of metabolic waste products unique to a high glucose environment and ROS may contribute to the observed decrease in cellular proliferation (25).

MPE and AM was used in respect of high level of antioxidants. The pericarps of *G. mangostana* L. is known for its high concentration of xanthenes and it has pharmacological effect as antioxidant (19). Under hyperglycaemic conditions, we suggest that antioxidants are able to regenerate a damaged ECM and improve cell growth as a result of oxidative stress through non-enzymatic glycation of proteins (32, 33). Because oxidative stress is associated with glomerulosclerosis and other disease related to a reduced antioxidant defense, therefore, it can be postulated that the

antioxidants, which can reduce the oxidative stress and prevent the progression of the disease, may exert a key role to protect mesangial cells in glomerulosclerosis (34, 35).

Present study shows, that MPE and AM increase cell proliferation and significantly reduced the level of TGF- β and fibronectin in glucose-induced mesangial cells compared to positive control. According to Jha *et al.* (2016) (36), antioxidants are able to convert ROS into nonreactive oxygen molecules which is harmless to cells (36). It also has an effect on retarding glucose absorption through inhibition of carbohydrate-hydrolyzing enzymes such as α -glucosidase and α -amylase (37) and down-regulates the TGF- β expression and fibronectin level (38) by decreasing NADPH oxidase expression. The expression of NADPH oxidase is elevated in diabetic nephropathy and it a source of oxidative stress. The up-regulation of NADPH oxidase subunits p47^{phox} and p22^{phox} plays an important role in ROS production and elevation fibronectin in high glucose condition (39). Antioxidant contained in MPE and AM may also ameliorate the antiproliferative response of mesangial cell to high level of glucose by altering gene transcription factors that act to regulate the cell growth (31).

According to Dennis & Witting (2017), anti-inflammatory agents may potentially reduce ROS via stabilizing endothelium function and NO bioactivity. Thus pathway may improve renal function and decrease tubular damage. Moreover, anti-inflammatory, as well as up-regulating gene responses, linked to antioxidant and cytoprotection (40). Down-regulation of TGF- β 1 and fibronectin level indicate an improvement of cell proliferation and metabolism in mesangial cells. This is presumably due to the influence of xanthone and α -mangostin contained in MPE which is able to neutralize free radicals that retarding cell damage (19, 23, 41).

Conclusion

Mangosteen peel extract (MPE) (in 5 and 20 μ g/ml) increased proliferation of cells in range glucose-induced concentration of 5-25 mM and significantly reduced TGF- β 1 and fibronectin levels in glucose-induced mesangial cells in glucose-induced concentration 5-10 mM. In conclusion, MPE products performed glucose induced-mesangial cells as *in vitro* model of diabetic glomerulosclerosis.

Acknowledgement

This research was funded and facilitated by Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, Indonesia. The author also thankful to Hanna Sari W Kusuma, Rismawati Laila Qodariah, Fajar Sukma Perdana, Annisa Arlisyah, Ni Luh Wisma Ekayanti from Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, Indonesia for their valuable assistance.

Conflict of Interest

The authors declare that no conflict of interest exists.

References

- Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010; 87:4-14.
- Kashihara N, Haruna Y, Kondeti VK, Kanwar YS. Oxidative stress in diabetic nephropathy. *Curr Med Chem* 2013; 17:4256-4269.
- Giacco F, Brownlee M. Oxidative stress and diabetic complications. *Circ Res* 2010; 107:1058-1070.
- Liu Y, Lu S, Zhang Y, Wang X, Kong F, Liu Y. Role of caveolae in high glucose and TGF- β 1 induced fibronectin production in rat mesangial cells. *Int J Clin Exp Pathol* 2014; 7:8381-8390.
- Qian Y, Feldman E, Pennathur S, Kretzler M, Brosius FC. From fibrosis to sclerosis mechanisms of glomerulosclerosis in diabetic nephropathy. *Diabetes* 2008; 57:1439-1445.
- Alsaad KO, Herzenberg AM. Distingusihing diabetic nephropathy from other causes of glomerulosclerosis: an update. *J Clin Pathol* 2007; 60:18-26.
- Zhu Y, Usui HK, Sharma K. Regulation of transforming growth factor-beta in diabetic nephropathy: implications for treatment. *Sem Nephrol* 2007; 27:153-160.
- Patil SB, Patil GS, Kundaragi VS, Biradar AN. A case of xanthogranulomatous pyelonephritis with spontaneous renocolic fistula. *Turk J Urol* 2013; 39:122-125.
- Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional food: impact on human health. *Pharmacog* 2010; 4:118-126.
- Chong YM, Chang SK, Sia WCM, Yim HS. Antioxidant efficacy of mangosteen (*Garcinia mangostana* Linn.) peel extracts in sunflower oil during accelerated storage. *Food Biosci.* 2015; 12:18-25.
- Pedraza-Chaccerri J, Cardenas-Rodriguez N, Orozco-Ibarra M, Perez-Rojas JM. Medicinal properties of mangosteen (*Garcinia mangostana*). *Food Chem Toxicol* 2008; 46:3227-3239.
- Tewtrakul S, Wattanapiromsakul C, Mahabusarakam W. Effects of compounds from *Garcinia mangostana* on inflammatory mediators in RAW 264.7 macrophage cells. *J Ethnopharm* 2009; 121:379-382.
- Widowati W, Darsono L, Suherman J, Fauziah N, Maesaroh M. Anti-inflammatory effect of mangosteen (*Garcinia mangostana* L.) peel extract and its compounds in LPS-induced RAW264.7 Cells. *Cells Nat Prod Sci* 2016; 22:147-153.
- Novilla A, Djamhuri DS, Fauziah N, Maesaroh M, Balqis B, Widowati W. Cytotoxic activity of mangosteen (*Garcinia mangostana* L.) peel extract and α -mangostin toward leukemia cell lines (HL-60 and K-562). *J Nat Remed* 2016; 16:52-59.
- Zadernowski R, Czaplicki S, Naczek M. Phenolic acid profiles of mangosteen fruits (*Garcinia mangostana*). *Food Chem* 2009; 112:153-160.
- Pothitirat W, Chomnawang MT, Supabphol R, Gritsanapan W. Comparison of 543 bioactive compounds content, free radical scavenging and anti-acne inducing bacteria 544 activities of extracts from the mangosteen fruit rind at two stages of 545 maturity. *Fito terapia* 2009; 80:422-447.
- Suttirak W, Manurakchinakom S. *In vitro* antioxidant properties of mangosteen peel extract. *J Food Sci Tech* 2012; 51:3546-3558.
- Widowati W, Rusmana D, Hardiman H, Tiono H, Wargasetia TL, Pujimulyani D. Mangosteen peel (*Garcinia mangostana* L.) extract for effervescent tablet. *Proceeding World Academy of Sci, Eng & Tech* 2013; 82:190-195.
- Widowati W, Darsono L, Suherman J, Yelliantty Y, Maesaroh M. High Performance Liquid Chromatography (HPLC) analysis, antioxidant, antiaggregation of mangosteen peel extract (*Garcinia mangostana* L.). *Int J Biosci Biochem Bioinform* 2014; 4:458-466.
- Adnyana IK, Abuzaid AS, Iskandar EY, Kurniati NF. Pancreatic lipase and α -amylase inhibitory potential of mangosteen (*Garcinia mangostana* Linn.) pericarp extract. *Int J Med Res Health Sci* 2016; 5:23-28.
- Abuzaid AS, Iskandar EY, Kurniati NF, Adnyana IK. Prevention of obesity and development of metabolic syndrome

- by mangosteen (*Garcinia mangostana* L.) pericarp ethanolic extract in male wistar rats fed with high-fat diet. *Int J Pharm Pharmaceut Sci* 2016; 2016; 8:1-7.
22. Abuzaid AS, Iskandar EY, Kurniati NF, Adanyana IK. Preventive effect on obesity of mangosteen (*Garcinia mangostana* L.) pericarp ethanolic extract by reduction of fatty acid synthase level in monosodium glutamate and high-calorie diet-induced male wistar rats. *Asian J Pharm Clin Res* 2016; 9:257-260.
23. Widowati W, Darsono L, Suherman J, Fauziah N, Maesaroh M, Erawijantari PP. Anti-inflammatory effect of mangosteen (*Garcinia mangostana* L.) peel extract and its compounds in LPS-induced RAW264.7 cells. *Nat Prod Sci* 2016; 22: 147-153.
24. Rusmana D, Elisabeth M, Widowati W, Fauziah N, Maesaroh M. Inhibition of inflammatory agent production by ethanolic extract and eugenol of *Syzygium aromaticum* flower bud (clove) in LPS-stimulated RAW264.7 cells. *Res J Med Plants* 2015; 9:264-274.
25. Nahman NS, Leonhart KL, Cosio FG, Hebert CL. Effects of high glucose on cellular proliferation and fibronectin production by cultured human mesangial cells. *Kidney Int* 1992; 41:396-402.
26. Dai MM, Wu H, Li H, Chen J, Chen JY, Hu SL, et al. Effects and mechanisms of Geniposide on rats with adjuvant arthritis. *Int Immunopharmacol* 2014; 20:46-53.
27. Pankov R, Yamada KM. Non-radioactive quantification of fibronectin matrix assembly. *Curr Protoc Cell Biol* 2004; 10:10-13.
28. Kolset SO, Reinbolt FP, Jenssen T. Diabetic nephropathy and extracellular matrix. *J Histochem Cytochem* 2012; 60:976-986.
29. Qian Y, Wang X, Liu Y, Li Y, Colvin RA, Tong L, et al. Extracellular ATP is internalized by macropinocytosis and induces intracellular ATP increase and drug resistance in cancer cells. *Cancer Lett* 2014; 351:242-251.
30. Loeffler I, Wolf G. Transforming growth factor- β and the progression of renal disease. *Nephrol Dial Transplant* 2013; 29:37-45.
31. Trachtman H, Futterweit S, Prenner J, Hanon S. Antioxidants reverse the antiproliferative effect of high glucose and advanced glycosylation end products in cultured rat mesangial cells. *Biochem Biophys Res Commun* 1994; 199: 346-352.
32. Chen C, Cheng K, Chang AY, Lin Y, Hseu Y, Wang H. 10-Shogaol, an Antioxidant from *Zingiber officinale* for skin cell proliferation and migration enhancer. *Int J Mol Sci* 2012; 13:1762-1777.
33. Kurtz A, Oh S. Age related changes of the extracellular matrix and stem cell maintenance. *Prev Med* 2012; 54:S50-S56.
34. Lee HS, Song CY. Oxidized low-density lipoprotein and oxidative stress in the development of glomerulosclerosis. *Am J Nephrol* 2009; 29:62-70.
35. David MS, Coombes JS, Bennett N, Johnson DW, Gobe GC. Oxidative stress, anti-oxidant therapies and chronic kidney disease. *Nephrology* 2012; 17:311-321.
36. Jha JC, Banal C, Chow BS, Cooper ME, Jandeleit-Dahm K. Diabetes and kidney disease: role of oxidative stress. *Antioxid Redox Signal* 2016; 25:657-684.
37. Manaharan T, Palanisamy UD, Ming CH. Tropical plant extract as potential antihyperglycemic agents. *Molecules* 2012; 17:5915-5923.
38. Han H, Cao A, Wang L, Guo H, Zang Y, Li Z. Huangqi decoction ameliorates streptozotocin-induced rat diabetic nephropathy through antioxidant and regulation of the TGF- β /MAPK / PPAR- γ signaling. *Cell Physiol Biochem* 2017; 42:1934-1944.
39. Zhang L, Pang S, Deng B, Qian L, Chen J, Zou J, et al. High glucose induces renal mesangial cell proliferation and fibronectin expression through JNK/NF- κ B/NADPH oxidase/ROS pathway, which is inhibited by resveratrol. *Int J Biochem Cell Biol* 2012; 44:629-638.
40. Dennis JM, Witting PK. Protective role for antioxidants in acute kidney disease. *Nutrients* 2017; 9:1-25.
41. Kurniawati M, Mahdi C, Aulanni'am A. The effect of juice mangosteen rind (*Garcinia mangostana* L.) to blood sugar levels and histological of pancreatic rats with the induction of streptozotocin. *J Pure App Chem Res* 2014; 3:1-6.